


Quantum anomalous Hall effect in intrinsic magnetic topological insulator

13 February 2020, by Thamarasee Jeewandara



Fabrication of MnBi₂Te₄ thin-flake device. (A) Optical image of representative few-layer MnBi₂Te₄ flakes cleaved onto Al₂O₃ thin film. The MnBi₂Te₄/Al₂O₃ stack is supported by a PDMS substrate. Image was obtained in transmission mode. Scale bar: 20 μm. (B) Optical image of the same MnBi₂Te₄/Al₂O₃ stack transferred on to a 285-nm SiO₂/Si substrate. Tape residue is visible under the Al₂O₃ film; the residue does not affect sample fabrication and subsequent measurements. (C) Optical image of a device fabricated from the sample shown in B. Metal (Cr/Au) contacts to the sample were thermally evaporated through a stencil mask. (D) Optical image of the same device after removing excess MnBi₂Te₄ flake that shorts adjacent electrodes with a sharp tip. Credit: *Science*, doi: 10.1126/science.aax8156


Nontrivial band topology can combine with magnetic order in a [magnetic topological insulator](#) to produce exotic states of matter such as [quantum anomalous Hall](#) (QAH) insulators and [axion insulators](#). An aim of condensed matter physics is to find new materials with useful properties and apply quantum mechanics to study them. The field has allowed physicists to better understand the uses of magnets for hard disk data storage, computer displays and other technologies.

The recent discovery of topological insulators have attracted broad interest and researchers predict that the interplay between ferromagnetism and the topological insulator state can realize a range of exotic quantum magnetic phenomena of interest in fundamental physics and device applications.

In a new report, Yujun Deng and a research team at the departments of physics and quantum matter physics in China, probed quantum transport in a thin flake MnBi₂Te₄ topological insulator, with intrinsic magnetic order. The ferromagnetic layers coupled anti-parallelly to each other in the atomically thin MnBi₂Te₄ layered van der Waals crystal. However, the sample became ferromagnetic when it contained an odd number of septuple layers. The research team observed the zero-field QAH effect in a five-septuple-layer specimen at 1.4 Kelvin. The results established MnBi₂Te₄ as an ideal platform to explore exotic topological phenomena with spontaneously broken [time-reversal symmetry](#). The work is now published on *Science*.

Topological materials distinctly contain [topologically protected quantum states](#) that are robust against local distresses. For instance, in a topological insulator (TI) such as [bismuth telluride](#) (Bi₂Te₃), the bulk band topology can guarantee the existence of two-dimensional (2-D) surface states with [gapless Dirac dispersion](#). By introducing magnetism into the initially time-reversal invariant topological insulators (TIs), scientists can induce profound changes in their electronic structure. For example, to experimentally observe the QAH effect in [chromium-doped \(Bi, Sb\)₂Te₃](#), physicists had to precisely control the ratio of multiple elements in a non-stoichiometric material. Fine-tuning the material required reconciling conflicting demands and therefore, researchers had to precisely quantize the anomalous Hall effect only at temperatures up to T = 2 K, far below the [Curie temperature](#) and [exchange gap in the material](#). To further explore the rich topological phenomena and their potential

applications, researchers must use intrinsic magnetic TIs (topological insulators) with an innate [magnetic order](#) to study their topological effects in pristine crystals.




Fabrication and characterization of few-layer MnBi_2Te_4 devices. (A) Optical image of few-layer flakes of MnBi_2Te_4 cleaved onto thermally-evaporated Al_2O_3 thin film (thickness ~ 70 nm). The $\text{MnBi}_2\text{Te}_4/\text{Al}_2\text{O}_3$ stack is supported on a PDMS substrate. Image was taken in transmission mode. Number of SLs are labeled on selected flakes. Scale bar: 20 μm . (B) Transmittance as a function of number of SLs. The transmittance (filled circles) follows the Beer-Lambert law (solid line). (C) Temperature-dependent sample resistance of few-layer MnBi_2Te_4 . The antiferromagnetic transition manifests as a resistance peak in the three-, four- and five-layer samples (Sample 3a, 4a and 5a, respectively; see table S1). Inset: layered crystal structure of MnBi_2Te_4 in the antiferromagnetic state. The spins of Mn^{2+} ions order ferromagnetically within a layer, whereas neighboring layers couple antiferromagnetically with an out-of-plane magnetocrystalline anisotropy. (D) R_{xy} of the same three-, four-, and five-layer MnBi_2Te_4 samples shown in C as a function of external magnetic field applied perpendicularly to sample plane. Data were obtained at $T = 1.6$ K. All data sets were anti-symmetrized to remove

R_{xx} component (23). The external magnetic field flips individual ferromagnetic SLs, one SL at a time, and eventually fully polarizes all SLs. The magnetic transitions manifest as jumps in R_{yx} that are marked by colored ticks on horizontal axes. Cartoons illustrate the magnetic states at representative magnetic fields (marked by open circles). SLs with up (down) magnetization are shown in red (blue). For simplicity, only one of the possible configurations are shown when there are degeneracies; we also ignore magnetic domains that may be present in some of the magnetic states. Credit: *Science*, doi: 10.1126/science.aax8156

In this work, Deng et al. probed quantum transport in atomically thin flakes of intrinsic magnetic topological insulator MnBi_2Te_4 . The material contained a layered [ternary tetradymite](#) compound containing septuple layers (Te-Bi-Te-Mn-Te-Bi-Te). The resulting MnBi_2Te_4 crystal was intrinsically magnetic and the magnetism originated from Mn^{2+} ions in the crystal. They studied thin flakes of MnBi_2Te_4 to minimize parallel bulk conduction and focused on MnBi_2Te_4 flakes containing an odd number of layers.


The team started with high quality MnBi_2Te_4 crystals grown using [a flux method](#) to obtain atomically thin MnBi_2Te_4 via [\$\text{Al}_2\text{O}_3\$ -assisted exfoliation](#). To accomplish this, they thermally evaporated Al_2O_3 thin film onto a freshly prepared surface of the bulk crystal, lifted the bulk using a thermal release tape then released the combined $\text{Al}_2\text{O}_3/\text{MnBi}_2\text{Te}_4$ stack on to a piece of transparent [polydimethylsiloxane](#) (PDMS) for microscopic inspection. Thereafter, they stamped the thin flakes onto a silicon wafer covered with SiO_2 , followed by deposition of Cr/Au contacts for transport measurements. The team completed the process in an airtight box to prevent sample exposure to oxygen (O_2) and water (H_2O) to mitigate sample degradation. They then extensively studied the rich set of magnetic states for the few-layered samples.



Quantum anomalous Hall effect in a five-layer MnBi₂Te₄ flake. (A and B) Magnetic-field-dependent Ryx (A) and Rxx (B) acquired in the five-layer Sample 5b at T = 1.4 K. Ryx and Rxx data shown here are anti-symmetrized and symmetrized, respectively, to remove the mixing of the two components (23). Up and down sweeps of the magnetic field are shown in red and blue, respectively. Ryx reaches 2 0.97 / h e , concomitant with a Rxx of 2 0.061 / h e at $\mu_0 H = 0$ T. These features are unambiguous evidence of zero-field QAH effect. External magnetic field polarizes the ferromagnetic SLs individually, and further improves the QAH quantization; Ryx quantizes to 2 0.998 / h e under magnetic fields above $\mu_0 H \sim 2.5$ T. (C) Rxx of Sample 5b as a function of magnetic field acquired at various temperatures. Data are symmetrized to remove Ryx component. (D) Arrhenius plot of Rxx as a function of 1/T under representative magnetic fields. Solid lines are line fits, the slope of which yields the energy gap of the thermally activated charge transport. (E) Energy gap as a function of magnetic field extracted from fitting the Arrhenius plots exemplified in D. Shaded region represents the error bound of the energy gap from the line fits. Solid circles highlight the representative gap values obtained from the fittings shown in D. Colored ticks on horizontal axes on panel B, C and E mark the location of magnetic transitions. All data were obtained under a backgate bias of $V_g = 200$ V. Credit: *Science*, doi: 10.1126/science.aax8156

Deng et al. observed a well-developed QAH effect at zero magnetic field in a five-layer MnBi₂Te₄ of much improved sample quality. They noted that an [external magnetic field](#) further improved the quantization by aligning the ferromagnetic layers. The ferromagnetic alignment also improved the robustness of the QAH effect against thermal fluctuations. At zero magnetic field, they obtained an energy gap that exceeded the value in [magnetically doped Ti thin films](#), although still much smaller than [the exchange gap expected](#) for MnBi₂Te₄.

The energy gap did not directly measure the bandgap of the surface states in the crystal, but characterized the minimum energy required to excite an electron from the valence to the conduction band. For instance, a large difference between the energy gap and the predicted bandgap implied various disorders in the sample. As a result, there is much room to further increase the energy scale of the QAH effect in pristine, high-quality MnBi₂Te₄ samples.



Gate-tuned quantum anomalous Hall effect in a five-layer MnBi₂Te₄ flake. (A) Magnetic-field-dependent Ryx, acquired in Sample 5b, under varying gate biases V_g (in 10 V steps). All data were obtained at T = 1.6 K. Curves are anti-symmetrized to remove Rxx component. Colored ticks on horizontal axis mark the location of magnetic

transitions. (B to D) R_{xx} and R_{yx} as functions of V_g under three representative magnetic fields, $\mu_0 H = 0$ T, 5 T and 14 T. An additional plateau of $\nu = 2$ emerges at $V_g \sim 25$ V, accompanied by a vanishing R_{xx} (panel D). The same plateau is also visible in A at $\mu_0 H > 10$ T during field sweeps under $V_g = 60$ V. This evidence points to a quantized Hall state with a filling factor $\nu = 2$. All data were obtained in the same Sample 5b, but V_g values do not exactly match those in A because of hysteresis during gate sweeps. Credit: *Science*, doi: 10.1126/science.aax8156

After the applied external magnetic field fully polarized the five-layer sample, the [energy gap](#) diminished with increasing magnetic field. The QAH states gradually evolved in the experimental setup providing a peek into the electronic structure of the surface bands outside the bandgap. Deng et al. understood all states observed in the study from a unified view. The Hall measurements near zero magnetic field yield a gate efficiency of 5×10^{10} cm^{-2}/V , which agreed well with the efficiency estimated from the device geometry. Since MnBi_2Te_4 is a layered material, the team expect the techniques developed for 2-D materials to be applicable to MnBi_2Te_4 . In this way, Yujun Deng and colleagues anticipate that van der Waals heterostructures integrating MnBi_2Te_4 with other magnetic/superconducting 2-D materials will provide fertile ground to further explore exotic topological quantum phenomena.

More information: Yujun Deng et al. Quantum anomalous Hall effect in intrinsic magnetic topological insulator MnBi_2Te_4 , *Science* (2020). [DOI: 10.1126/science.aax8156](https://doi.org/10.1126/science.aax8156)

Haijun Zhang et al. Topological insulators in Bi_2Se_3 , Bi_2Te_3 and Sb_2Te_3 with a single Dirac cone on the surface, *Nature Physics* (2009). [DOI: 10.1038/nphys1270](https://doi.org/10.1038/nphys1270)

Yoshinori Tokura et al. Magnetic topological insulators, *Nature Reviews Physics* (2019). [DOI: 10.1038/s42254-018-0011-5](https://doi.org/10.1038/s42254-018-0011-5)

© 2020 Science X Network

APA citation: Quantum anomalous Hall effect in intrinsic magnetic topological insulator (2020, February

13) retrieved 17 January 2021 from <https://phys.org/news/2020-02-quantum-anomalous-hall-effect-intrinsic.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.