


# How newborn stars prepare for the birth of planets

20 February 2020



ALMA and the VLA observed more than 300 protostars and their young protoplanetary disks in Orion. This image shows a subset of stars, including a few binaries. The ALMA and VLA data compliment each other: ALMA sees the outer disk structure (visualized in blue), and the VLA observes the inner disks and star cores (orange). Credit: ALMA (ESO/NAOJ/NRAO), J. Tobin; NRAO/AUI/NSF, S. Dagnello

An international team of astronomers used two of the most powerful radio telescopes in the world to create more than three hundred images of planet-forming disks around very young stars in the Orion Clouds. These images reveal new details about the birthplaces of planets and the earliest stages of star formation.


Most of the [stars](#) in the universe are accompanied by planets. These planets are born in rings of dust and gas, called protoplanetary disks. Even very young stars are surrounded by these disks. Astronomers want to know exactly when these disks start to form, and what they look like. But young stars are very faint, and there are dense [clouds](#) of dust and gas surrounding them in stellar nurseries. Only highly sensitive radio telescope arrays can spot the tiny disks around these infant

stars amidst the densely packed material in these clouds.

For this new research, astronomers pointed both the National Science Foundation's Karl G. Jansky Very Large Array (VLA) and the Atacama Large Millimeter/submillimeter Array (ALMA) to a region in space where many stars are born: the Orion Molecular Clouds. This survey, called VLA/ALMA Nascent Disk and Multiplicity (VANDAM), is the largest survey of young stars and their disks to date.


Very young stars, also called protostars, form in clouds of gas and dust in space. The first step in the formation of a star is when these dense clouds collapse due to gravity. As the cloud collapses, it begins to spin—forming a flattened [disk](#) around the [protostar](#). Material from the disk continues to feed the star and make it grow. Eventually, the left-over material in the disk is expected to form planets.

Many aspects about these first stages of star formation, and how the disk forms, are still unclear. But this new survey provides some missing clues as the VLA and ALMA peered through the dense clouds and observed hundreds of protostars and their disks in various stages of their formation.



### Four special protostars

Among hundreds of survey images, four protostars looked different than the rest and caught the scientists' attention. "These newborn stars looked very irregular and blobby," said team member Nicole Karnath of the University of Toledo, Ohio (now at SOFIA Science Center). "We think that they are in one of the earliest stages of star formation and some may not even have formed into protostars yet."



The Orion Molecular Clouds, the target of the VANDAM survey. Yellow dots are the locations of the observed protostars on a blue background image made by Herschel. Side panels show nine young protostars imaged by ALMA (blue) and the VLA (orange). Credit: ALMA (ESO/NAOJ/NRAO), J. Tobin; NRAO/AUI/NSF, S. Dagnello; Herschel/ESA

Schematic shows a proposed pathway (top row) for the formation of protostars, based on four very young protostars (bottom row) observed by VLA (orange) and ALMA (blue). Step 1 represents the collapsing fragment of gas and dust. In step 2, an opaque region starts to form in the cloud. In step 3, a hydrostatic core starts to form due to an increase in pressure and temperature, surrounded by a disk-like structure and the beginning of an outflow. Step 4 depicts the formation of a class 0 protostar inside the opaque region, that may have a rotationally supported disk and more well-defined outflows. Step 5 is a typical class 0 protostar with outflows that have broken through the envelope (making it optically visible), an actively accreting, rotationally supported disk. In the bottom row, white contours are the protostar outflows as seen with ALMA. Credit: ALMA (ESO/NAOJ/NRAO), N. Karnath; NRAO/AUI/NSF, B. Saxton and S. Dagnello

### Young planet-forming disks

"This survey revealed the average mass and size of these very young [protoplanetary disks](#)," said John Tobin of the National Radio Astronomy Observatory (NRAO) in Charlottesville, Virginia, and leader of the survey team. "We can now compare them to older disks that have been studied intensively with ALMA as well."

What Tobin and his team found, is that very young disks can be similar in size, but are on average much more massive than older disks. "When a star grows, it eats away more and more material from the disk. This means that younger disks have a lot more raw material from which planets could form. Possibly bigger planets already start to form around very [young stars](#)."

It is special that the scientists found four of these objects. "We rarely find more than one such irregular object in one observation," added Karnath, who used these four infant stars to propose a schematic pathway for the earliest stages of star

formation. "We are not entirely sure how old they are, but they are probably younger than ten thousand years."

To be defined as a typical (class 0) protostar, stars should not only have a flattened rotating disk surrounding them, but also an outflow—spewing away material in opposite directions—that clears the dense cloud surrounding the stars and makes them optically visible. This outflow is important, because it prevents stars from spinning out of control while they grow. But when exactly these outflows start to happen, is an open question in astronomy.

One of the infant stars in this study, called HOPS 404, has an outflow of only two kilometers (1.2 miles) per second (a typical protostar-outflow of 10-100 km/s or 6-62 miles/s). "It is a big puffy sun that is still gathering a lot of mass, but just started its outflow to lose angular momentum to be able to keep growing," explained Karnath. "This is one of the smallest outflows that we have seen and it supports our theory of what the first step in forming a protostar looks like."

### Combining ALMA and VLA

The exquisite resolution and sensitivity provided by both ALMA and the VLA were crucial to understand both the outer and inner regions of protostars and their disks in this survey. While ALMA can examine the dense dusty material around protostars in great detail, the images from the VLA made at longer wavelengths were essential to understand the inner structures of the youngest protostars at scales smaller than our solar system.

"The combined use of ALMA and the VLA has given us the best of both worlds," said Tobin. "Thanks to these telescopes, we start to understand how planet formation begins."

**More information:** John J. Tobin et al, The VLA/ALMA Nascent Disk and Multiplicity (VANDAM) Survey of Orion Protostars. II. A Statistical Characterization of Class 0 and Class I Protostellar Disks, *The Astrophysical Journal* (2020). DOI: [10.3847/1538-4357/ab6f64](https://doi.org/10.3847/1538-4357/ab6f64)

N. Karnath et al. Detection of Irregular,

Submillimeter Opaque Structures in the Orion Molecular Clouds: Protostars within 10,000 yr of Formation?, *The Astrophysical Journal* (2020). DOI: [10.3847/1538-4357/ab659e](https://doi.org/10.3847/1538-4357/ab659e)

Provided by National Radio Astronomy Observatory

APA citation: How newborn stars prepare for the birth of planets (2020, February 20) retrieved 21 January 2021 from <https://phys.org/news/2020-02-newborn-stars-birth-planets.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*