



# Breakthrough made toward more powerful particle accelerators

11 March 2020, by Joohyeon Heo

## MICE Muon Ionization Cooling Experiment

MICE has made the first ever demonstration of the ionization cooling of muons – a major step in the journey to create the world's most powerful particle accelerator.

- 1 Bunches of protons are accelerated into a target of dense material (such as tungsten or mercury). The atoms within the target emit a particle called a pion.
- 2 Pions are unstable and they quickly decay into a muon and a neutrino.
- 3 The neutrinos, being virtually massless and without charge, pass out of the experiment. Magnets direct charged muons of the correct energy moving in the right direction.



Infographic: STFC, Ben Gilliland


Muon ionization cooling experiment by the MICE collaboration. Credit: MICE collaboration

An international team of researchers, affiliated with UNIST has for the first time demonstrated the ionization cooling of muons. Regarded as a major step in creating more powerful particle accelerators, this new muon accelerator is expected to provide a better understanding of the fundamental constituents of matter.

This breakthrough has been carried out by the Muon Ionization Cooling Experiment (MICE) collaboration, which includes many U.K. scientists, as well as Professor Moses Chung and his research team in the School of Natural Sciences at UNIST. Their findings were published in the online version of *Nature* on February 5, 2020.

"We have succeeded in realizing muon ionization cooling, one of our greatest challenges associated with developing muon accelerators," says Professor Chung. "This achievement is considered especially important, as it could change the paradigm of developing the lepton collider that could replace the Neutrino Factory or the Large Hadron Collider (LHC)."

Muons are naturally occurring particles generated in the Earth's upper atmosphere by cosmic ray collisions, and are thus regarded as a follow-on particle accelerator to replace the LHC. Protons, a type of hadron, are primarily used by the LHC and they partake in strong interactions. Leptons, like the electron and the muon, are not subject to the strong interaction; rather, they interact via the weak force.



The MICE muon beam-line at the Science and Technology Facilities Council (STFC) ISIS Neutron and Muon Beam facility on the Harwell Campus in the UK.

Credit: MICE collaboration

Muons have an extremely brief lifespan of two-millionths of a second. They are produced by smashing a [beam](#) of protons into a target. These muons form a diffuse cloud, meaning that they are difficult to accelerate and there is a low chance of them colliding and producing useful physical phenomena. To make the cloud less diffuse, a process known as beam cooling was suggested. This involves getting the muons closer together and moving in the same direction. However, due to the ultra-short lifespan of muons, it has been impossible to cool the beam using traditional methods.

To tackle this challenge, the MICE collaboration team succeeded in channeling muons into a small enough volume via a method known as ionization cooling, which was previously suggested and developed into theoretically operable schemes in the 1980s.

The results of the experiment, carried out using the MICE muon beamline at the Science and Technology Facilities Council (STFC) ISIS Neutron and Muon Beam facility on the Harwell Campus in the U.K., clearly shows that the phase-space volume occupied by the [muon](#) beam can be controlled via [ionization](#) cooling, as predicted by the theory.

**More information:** Demonstration of cooling by the Muon Ionization Cooling Experiment, *Nature* (2020). [DOI: 10.1038/s41586-020-1958-9](https://doi.org/10.1038/s41586-020-1958-9)

Provided by Ulsan National Institute of Science and Technology

APA citation: Breakthrough made toward more powerful particle accelerators (2020, March 11) retrieved 20 January 2021 from <https://phys.org/news/2020-03-breakthrough-powerful-particle.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*