

India's Chandrayaan 2 is creating the highest-resolution map we have of the moon

12 March 2020, by Jatan Mehta



Credit: ISRO

India's space organization, ISRO, launched Chandrayaan 2 to the moon last year in July. While its lander Vikram crashed on the lunar surface on September 7, the Chandrayaan 2 orbiter continues to orbit the moon.

The Chandrayaan 2 orbiter hosts an extensive set of instruments to map the [moon](#), and now, we get a peek at the data it has sent.

ISRO scientists had submitted a raft of initial results from the orbiter's mapping instruments to present at the flagship 51st Lunar and Planetary Science Conference in March. This is an annual conference hosted in the United States where more than 2000 planetary scientists and students from around the world attend and present their latest work. However, due to concerns about the novel coronavirus, the conference has been


anceled.

Seeing a crater in the dark

Chandrayaan 2 orbiter has an optical camera called the Orbiter High-Resolution Camera (OHRC) that captures detailed images of the moon. OHRC can image at a best resolution of 0.25 meters/pixel, beating NASA Lunar Reconnaissance Orbiter's (LRO) best of 0.5 meters/pixel.

Back in October, we already saw OHRC flex its muscles by sending images including clearly visible boulders less than 1 meter in size. And now OHRC has demonstrated imaging an area not directly illuminated by sunlight. It captured an image of a crater floor in shadow by seeing the dim light falling on it that has been reflected from the crater rim.

Moving ahead, this capability will be used to image insides of craters on the lunar poles, where sunlight never reaches. Mapping the terrain of polar craters is important because future lunar habitats are believed to be stationed near them, transporting [water](#) and other resources from inside them.



The Chandrayaan 2 orbiter. Credit: ISRO


Highest resolution 3-D maps

The Terrain Mapping Camera (TMC 2) onboard Chandrayaan 2 is a stereo imager, meaning it can capture 3-D images. It does that by imaging the same site from three different angles, akin to NASA's LRO, from which a 3-D image is constructed.

TMC 2 has beamed back images taken from 100 km above the [lunar surface](#) and the 3-D views generated from them look great. Here is one of a crater and a wrinkled ridge, the latter being a tectonic feature.

Such images are very useful for understanding how lunar features form and get their shape. For example, a 3-D image can help construct an accurate picture of the geometry of the impact that formed a crater.

Over time, Chandrayaan 2 will provide the highest resolution 3-D images of the entire moon, the best case resolution being 5 meters/pixel.



Left: Image of the lunar surface by the Chandrayaan 2 orbiter. Region R1 is part of a crater not receiving sunlight at the time of image capture. Right: Crater floor in the dark imaged by Chandrayaan 2's OHRC by seeing the dim reflected light from the crater's rim. Credit: ISRO

Enhanced eyes in the infrared

The Imaging Infrared Spectrometer (IIRS) on

Chandrayaan 2 is the successor to the famous Moon Mineralogical Mapper (M3) instrument onboard Chandrayaan 1.


The M3 instrument, which was contributed by NASA, has been publicly acknowledged for its excellent mineral mapping capabilities and detection of water on the moon. Noah Petro, project scientist for LRO, recently noted on Twitter:

10 years ago today Chandrayaan-1 ended. I was so lucky to be a small part of that mission. The M3 instrument allowed up to take a huge step forward in learning about the composition of our 8th continent.

[@_DanOnTheMoon](#) [@pyroxena](#)
[@NASAmoon](#) [@isro](#)


— Noah Petro (@nepetro) [August 28, 2019](#)

Both IIRS and M3 detect reflected sunlight from the moon's surface. Scientists identify minerals on the surface based on the patterns of these reflections. The IIRS boasts nearly twice the sensitivity of M3 in infrared light and the initial results demonstrate to that effect.



3-D view of a crater on the moon generated from images captured by Chandrayaan 2 orbiter's Terrain Mapping Camera. Credit: ISRO

Thanks to M3, scientists now know that the lunar soil does hold trace amounts of water and hydroxyl molecules even in non-polar regions. IIRS onboard Chandrayaan 2 will map water concentrations in the lunar soil with improved sensitivity. Chandrayaan 2's long-term observations aim to discern how the water content in the lunar soil changes in response to the lunar environment, i.e., what the lunar water cycle looks like.




Note that all this is still less amount of water than the driest deserts on Earth. However, the [lunar poles](#) host appreciably more water. And that is where Chandrayaan 2's radar comes into the picture.

Quantifying water on the moon


The Dual Frequency Synthetic Aperture Radar (DFSAR) onboard the Chandrayaan 2 orbiter is the successor to the Miniature Synthetic Aperture Radar (Mini-SAR) on Chandrayaan 1. DFSAR penetrates the moon's surface twice as deeply as Mini-SAR. Not just that, DFSAR also boasts a higher resolution than the radar onboard LRO called Mini-RF. The initial results demonstrate as much, comparing a DFSAR radar image of region with Mini-RF.

3-D view of a wrinkled ridge on the moon generated from images captured by Chandrayaan 2 orbiter's Terrain Mapping Camera. Credit: ISRO

With greater penetration depth and higher resolution than any prior instruments, Chandrayaan 2's orbiter is in the process of adequately quantifying just how much water ice is trapped beneath the permanently dark crater floors on the moon's poles. Current estimates based on past observations suggest that the moon's poles host more than 600 billion kg of water ice, equivalent to at least 240,000 Olympic-sized swimming pools.



The Glauber crater on the moon imaged in infrared by Chandrayaan 2's IIRS and Chandrayaan 1's M3 respectively. Credit: ISRO, NASA



A region on the moon imaged by ISRO Chandrayaan 2's radar (leftmost), NASA LRO's radar (center) and LRO's visible light camera. Credit: ISRO

What's next?

The lunar science and exploration communities agree that we can harness water ice on the moon's poles to power future lunar habitats. Using solar power generated by the habitats, we can also split the water ice into hydrogen and oxygen for use as rocket fuel.

But before we plan habitats at the moon's poles, we need to know more about the nature of water ice in these regions and how to access it given their terrain. The initial results from Chandrayaan 2 clearly show the promise of the highest resolution mapper ever sent to the moon. ISRO has stated that Chandrayaan 2 will orbit the moon for seven years and that should be ample time to fully map and quantify water and their host regions on the moon.

Surface missions that explore these water-hosting permanently shadowed regions, like NASA's upcoming VIPER rover, are the next logical step towards sustainable habitats on the moon. As we develop technologies that tap into water ice on the moon, we can colonize not just our celestial neighbor but the solar system. We should be glad our moon has plenty of water; we can't keep dragging everything out of Earth's gravitational well forever.

Provided by Universe Today

APA citation: India's Chandrayaan 2 is creating the highest-resolution map we have of the moon (2020, March 12) retrieved 30 March 2020 from <https://phys.org/news/2020-03-india-chandrayaan-highest-resolution-moon.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.