

Small horses got smaller, big tapirs got bigger 47 million years ago

24 March 2020



Window into the 47 million year old ecosystem of the Geiseltal fossil locality with the small-sized horse-ancestor Propalaeotherium on the left, the ancient tapir Lophiodon in the middle, and a young terrestrial crocodile Bergisuchus in the background. Credit: Márton Szabó


The former coalfield of Geiseltal in eastern Germany has yielded large numbers of exceptionally preserved fossil animals, giving palaeontologists a unique window into the evolution of mammals 47 million years ago. At that time—the middle Eocene—the Earth was much warmer and the area was a swampy subtropical forest whose inhabitants included ancestors of the horse, ancient tapirs, large terrestrial crocodiles, as well as giant tortoises, lizards and ground-dwelling birds. So rich are the Geiseltal finds that they give researchers an unprecedented high-resolution picture of evolutionary dynamics at the population level.

A team led by Dr. Márton Rabi from the University of Tübingen and the Martin Luther University Halle-Wittenberg (MLU) has shown that the [body](#) size of two species of mammals developed in opposite directions. The study, published in *Scientific Reports*, was carried out with Simon Ring and

Professor Hervé Bocherens at the Senckenberg Centre for Human Evolution and Palaeoenvironment and the University of Tübingen in cooperation with Dr. Oliver Wings from the MLU.

"We were initially interested in the evolution of the ancient horses, which were about the size of a Labrador dog. These animals are particularly abundant in the Geiseltal fossil record," Rabi says. Researchers initially believed they had several species of early horse. "However, we found that here, there was only one species, whose body size shrank significantly with time," Rabi explains. The team wanted to test whether this body size shift was climate-induced, since past global warming caused body-size reduction in ancient mammals.

Carbon and oxygen isotope studies on fossil teeth provided the scientists with information about the local middle Eocene climate. "They indicate a humid tropical climate. However, we didn't find any evidence for [climatic changes](#) in Geiseltal over the period investigated," says Bocherens. To further test the data, the team sought to discover whether the dwarfing process was unique to the horses. For comparison, they examined the evolution of the tapir ancestor called Lophiodon. "We had reason to question the Geiseltal's constant-climate data; so we expected that other mammals would show the same body-size trends as the horses," Simon Ring explains. In a surprising result, the tapirs—also a single species—revealed the opposite trend. They grew larger instead of shrinking. While the ancestors of the horse shrank from an average body weight of 39 kilograms to around 26 kilograms over about a million years, the tapirs increased from 124 kilograms to an average body weight of 223 kilograms.



extends their lives and gives them more time to breed. The Geiseltal tapirs and the horses therefore likely maximized the different advantages of their respective life cycle strategies, which caused divergent [body size](#) evolution.

Exceptional fossil deposits

The Geiseltal fossil site is located in the eastern state of Saxony-Anhalt. In the course of open-cast brown coal mining between 1933 and 1993, tens of thousands of fossil specimens of more than one hundred species were discovered there. Many were the ancestors of modern vertebrates. "The Geiseltal is as important a fossil site as the Messel Pit near Darmstadt, which is a UNESCO World Heritage Site," says Dr. Rabi. "But because the Geiseltal collection was hardly accessible during East German times, it kind of went off the radar."

More information: Simon J. Ring et al. Divergent mammalian body size in a stable Eocene greenhouse climate, *Scientific Reports* (2020). [DOI: 10.1038/s41598-020-60379-7](https://doi.org/10.1038/s41598-020-60379-7)

Provided by Universitaet Tübingen

Exceptionally well fossilized skeletons of the ancient tapir *Lophiodon* (top) and the ancestral horse *Propalaeotherium* (bottom) from the middle Eocene Geiseltal locality (Germany, Saxony-Anhalt). Credit: Oliver Wings

Differing survival strategies

"All the data indicate that the body size of the horses and tapirs developed differently not because of the climate, but because of different life cycles," explains Bocherens. Small animals reproduce faster and die younger: Relative to their size, they don't have to eat as much to maintain their body mass and can devote more resources to having young. Larger animals live longer and have lower reproduction rates. They have to eat more and therefore have fewer resources for reproduction—but, being large, face fewer predators and can range further to get better food. That

APA citation: Small horses got smaller, big tapirs got bigger 47 million years ago (2020, March 24)
retrieved 26 February 2021 from <https://phys.org/news/2020-03-small-horses-smaller-big-tapirs.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.