


# International ozone treaty stops changes in Southern Hemisphere winds

March 25 2020

---



The 2019 ozone hole reached a peak extent of 6.3 million square miles on September 8, 2019, the lowest maximum observed in decades. This NASA visualization depicts ozone concentrations in Dobson Units, the standard measure for stratospheric ozone. Credit: NASA

Chemicals that deplete Earth's protective ozone layer have also been triggering changes in Southern Hemisphere atmospheric circulation. Now, new research in *Nature* finds that those changes have paused and

might even be reversing because of the Montreal Protocol, an international treaty that successfully phased out use of ozone-depleting chemicals.

"This study adds to growing evidence showing the profound effectiveness of the Montreal Protocol. Not only has the treaty spurred healing of the [ozone layer](#), it's also driving recent changes in Southern Hemisphere air circulation patterns," said lead author Antara Banerjee, a CIRES Visiting Fellow at the University of Colorado Boulder who works in the Chemical Sciences Division of the National Oceanic and Atmospheric Administration (NOAA). She started this work as a Postdoctoral Fellow at Columbia University.

The ozone hole, discovered in 1985, has been forming every spring in the atmosphere high over Antarctica. Ozone depletion cools the air, strengthening the winds of the polar vortex and affecting winds all the way down to the lowest layer of Earth's atmosphere. Ultimately, ozone depletion has shifted the midlatitude jet stream and the dry regions at the edge of the tropics toward the South Pole.

Previous studies have linked these circulation trends to weather changes in the Southern Hemisphere, especially rainfall over South America, East Africa, and Australia, and to changes in ocean currents and salinity.

The Montreal Protocol of 1987 phased out production of ozone-destroying substances such as chlorofluorocarbons (CFCs). Beginning around 2000, concentrations of those chemicals in the stratosphere started to decline and the ozone hole began to recover. In this study, Banerjee and her co-authors have shown that around the year 2000, the circulation of the Southern Hemisphere also stopped expanding polewards—a pause or slight reversal of the earlier trends.

"The challenge in this study was proving our hypothesis that ozone

recovery is in fact driving these [atmospheric circulation](#) changes and it isn't just a coincidence," Banerjee said.



This panoramic image from February 14, 2014, shows parts of Chile and Argentina from the International Space Station (ISS). Tierra del Fuego and Cape Horn are shown here, with the Atlantic Ocean in the foreground and the Pacific Ocean across the top of the image. Credit: NASA

To do that, the researchers used a two-step statistical technique called detection and attribution: detecting whether certain patterns of observed wind changes are unlikely to be due to natural variability alone and, if so, whether the changes can be attributed to human-caused factors, such as emissions of ozone-depleting chemicals and CO<sub>2</sub>.

Using computer simulations, the researchers first determined that the observed pause in circulation trends couldn't be explained by natural shifts in winds alone. Next, they isolated the effects of ozone and greenhouse gases separately.

They showed that while rising CO<sub>2</sub> emissions have continued expanding the near-surface circulation (including the jet stream) polewards, only the ozone changes could explain the pause in circulation trends. Prior to 2000, both [ozone depletion](#) and rising CO<sub>2</sub> levels pushed the near-surface circulation poleward. Since 2000, CO<sub>2</sub> has continued to push this circulation poleward, balancing the opposing effect of the ozone recovery.

"Identifying the ozone-driven pause in circulation trends in real-world observations confirms, for the first time, what the scientific ozone community has long predicted from theory," said John Fyfe, a scientist at Environment and Climate Change Canada and one of the paper's co-authors.

With ozone beginning to recover and CO<sub>2</sub> levels continuing to climb, the future is less certain, including for those Southern Hemisphere regions whose weather is affected by the jet stream and those at the edge of the dry regions.

"We term this a 'pause' because the poleward [circulation](#) trends might resume, stay flat, or reverse," Banerjee said. "It's the tug of war between the opposing effects of [ozone](#) recovery and rising greenhouse gases that will determine future trends."

**More information:** A pause in Southern Hemisphere circulation trends due to the Montreal Protocol, *Nature*, [DOI: 10.1038/s41586-020-2120-4](#) , [nature.com/articles/s41586-020-2120-4](#)

Provided by University of Colorado at Boulder

Citation: International ozone treaty stops changes in Southern Hemisphere winds (2020, March 25) retrieved 19 September 2024 from <https://phys.org/news/2020-03-international-ozone-treaty-southern-hemisphere.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.