

Overcoming carbon loss from farming in peatlands

March 31 2020, by Kaine Korzekwa



In this field with histosol soil, miscanthus was added as a form of organic matter. It is a type of biomass crop they studied to see if it can add carbon to the soil.
Credit: Jacynthe Dessureault-Rompré

In many regions of the world, farming must be done on areas of soil categorized as histosols. Histosols have a thick layer of rich organic matter, called peat.

Scientists are concerned, because farming can cause these soils to lose valuable carbon.

That's where Jacynthe Dessureault-Rompré and her team at Laval University in Canada come in. She is trying to show that histosols can be sustainably used for farming.

To do this, the research team performed a two-step experiment that involved adding different kinds of [plant material](#) to the soil. Their work is similar to how a backyard gardener may add compost to soil to add nutrients that have been lost.

"The first objective was to get a better understanding of plant material decomposition," Dessureault-Rompré says. "We looked at decomposition qualities of specific plant materials and how these affected long-term carbon stores in the soil. The second objective was to determine which plant performed best based on simulations of the long-term soil carbon storage."

Carbon in these soils is lost by erosion, tillage and a [natural process](#) called mineralization. The carbon is released from the soil as [carbon dioxide](#), a harmful greenhouse gas, into the atmosphere.



This field of histosols has willow on top that is ready to be incorporated so the researchers can study how it decomposes. Credit: Jacynthe Dessureault-Rompré

For their first experiment, the team used three common biomass crops: sorghum, miscanthus, and willow. They put plant material from these crops inside permeable bags and placed the bags directly in the soil.

Then, they analyzed which plant materials had the best decomposition characteristics. The ones that broke down slowly are the best for storing carbon in the soil longer.

"The field decomposition study gave us data on what happened to the

three different plant materials over a period of 17 months," Dessureault-Rompré says. "A good candidate is a crop that will last longer in the soil because the buildup of the carbon stock will be more efficient, therefore you need less biomass applied each year. The stability characteristics of the plant material are very important."

Next, researchers used decomposition data to simulate how much each plant would help the soil over a long period of time. They found that miscanthus and willow performed much better than sorghum. They also calculated amounts of the [plants](#) that would help the soils be the most sustainable.

"If you have a crop such as miscanthus that decomposes less than sorghum, the buildup over the years is much more efficient," she explains. "The simulation part added a new perspective because we were then able to see that carbon equilibrium is something that can be achieved. It was fantastic to see that adding plant material year after year allows farmers to overcome the [carbon](#) lost during farming in histosols."



The researchers obtained the biomass crops from fields before adding it to the histosols. These are miscanthus flowers. Credit: Jacynthe Dessureault-Rompré

She adds that it is hard to estimate when these farmers could adopt this practice. However, it's possible that within the next 10 years, this new soil conservation practice could be used by farmers.

While many scientists don't think histosols should be used for farming, many of the farmers have no choice. For the agricultural community, processing facilities and distribution services to make a living, farmers must grow crops on the land available to them.

The findings from Dessureault-Rompré and her team are important for easing concerns. This research demonstrates that it is possible to farm these soils sustainably.

"I was afraid this research would be criticized because it is a very new way of looking at crop production on these very special soils," she says. "Many scientists believe that cultivated histosols should be brought back to their natural state or that the degradation process of these [soil](#) is irreversible. But this project really aims to develop a sustainable way of growing high value [crops](#) on these soils.

More information: Jacynthe Dessureault-Rompré et al. Biomass crops as a soil amendment in cultivated histosols: can we reach carbon equilibrium?, *Soil Science Society of America Journal* (2020). [DOI: 10.1002/saj2.20051](#)

Provided by American Society of Agronomy

Citation: Overcoming carbon loss from farming in peatlands (2020, March 31) retrieved 26 April 2024 from <https://phys.org/news/2020-03-carbon-loss-farming-peatlands.html>

| |
|--|
| <p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p> |
|--|