


Seeking 'soundwaves' in the superfluid order parameter

April 14 2020



Ultracold atomic lab at Swinburne University of Technology. Credit: FLEET

A Swinburne University of Technology study published this week examines the propagation of energy as sound waves in a quantum gas, revealing for the first time strong variations in the nature of the sound wave as a function of temperature.

At low energies, this energy travels via the collective movement of many particles moving in sync—essentially, as sound waves—quantified using quasiparticles known as phonons.

Below the superfluid transition temperature T_c these sound waves in a unitary Fermi gas can propagate without collisions and are driven by

ripples in the phase of the superfluid order parameter (wavefunction)—this mode is known as the Bogoliubov-Anderson (BA) phonon.

Above T_c , the [sound waves](#) become more strongly-damped, and collisions play a dominant role.

- lower than T_c In the colder, superfluid mode, damping is dominated by collisions with thermally excited quasiparticles and is well described by (QRPA) theory
- $>T_c$ Above the transition temperature, the strongly-damped mode occurs at the crossover between collisionless hydrodynamic regimes.
- $\gg T_c$ At even [higher temperatures](#), collective propagation of sound-wave vanishes, and excitation is dominated by the energy of individual particles.

Strong similarities were identified in the temperature dependence of sound in the unitary Fermi gas and the behaviour of phonons in liquid helium, which was one of the first superfluids identified historically.

This study provides quantitative benchmarks for dynamical theories of strongly-correlated fermions.



The ultracold atomic gases formed and studied in Prof Chris Vale's lab at Swinburne allow very precise tuning of interactions between atoms.

"We cooled and confined a highly dilute gas of Li⁶ atoms, realising a unitary Fermi gas, which exhibits the strongest interactions allowed by quantum mechanics with a contact potential," explains Prof Vale.

In a unitary gas, elastic collisions become resonant and the thermodynamic properties of the gas become universal functions of the

temperature and density. Unitary Fermi gases allow precise testing of theories of interacting fermions.

The team then studied excitations in the gas above and below the superfluid phase transition T_c using two-photon Bragg spectroscopy.



Above: Excitation spectra for unitary Fermi gas showing (top) experimental data and (b) theory. Credit: FLEET


"We measured excitation spectra at a momentum of approximately half the Fermi momentum, both above and below the superfluid critical temperature T_c ," explains study author Dr. Carlos Kuhn.

Two, focused [laser pulses](#) (approx 1.2 milliseconds in duration) intersecting within the gas create a periodic perturbation for the lithium atoms.

Immediately after the twin laser pulse, the confining optical trap is switched off and the momentum of atoms is measured after 4 milliseconds of expansion, and can be mapped as a function of laser frequency.

The finite duration and size of the Bragg beams lead to a Fourier-limited spectral resolution of approximately 1:25 kHz FWHM which is well below the typical Fermi energies, EF 11 kHz, used in the experiments.

"High Frequency Sound in a Unitary Fermi Gas" was published in *Physical Review Letters* (PRL) in March 2020.



Ultracold atomic lab at Swinburne University of Technology. Credit: FLEET

Ultracold studies at FLEET

The study of many-body quantum systems with strong inter-particle interactions is of great interest for the understanding of novel materials.

Within FLEET, Chris Vale studies topological phenomena in 2-D gases of ultracold fermionic atoms, investigating cold atom implementations of Floquet topological superfluidity, nonequilibrium enhancements to the superconducting critical temperature and new forms of topological matter based on optically induced [spin-orbit coupling](#) in 2-D atomic gases, in Research Theme 3.

FLEET's research theme 3 studies systems that are temporarily driven out of thermal equilibrium to investigate the qualitatively different physics displayed and new capabilities for dynamically controlling their behaviour.

Chris leads the study of quantum gases at Swinburne University of Technology. In these collections of atoms cooled to only 100 nano-Kelvins above Absolute Zero, behaviours that are usually only found at the microscopic level become prominent at the macroscopic level.

The team's study of Fermi gases confined to 2-D tests new paradigms for dissipationless transport in topological and non-equilibrium quantum matter synthesised from ultracold atoms.

Chris is one of almost 100 researchers at FLEET, all motivated by one grand challenge: to reduce the energy used in information and communication technology (ICT), which already accounts for at least 8% of global electricity use, and is doubling every decade.

More information: C. C. N. Kuhn et al. High-Frequency Sound in a Unitary Fermi Gas, *Physical Review Letters* (2020). [DOI: 10.1103/physrevlett.124.150401](#)

Provided by FLEET

Citation: Seeking 'soundwaves' in the superfluid order parameter (2020, April 14) retrieved 25 April 2024 from <https://phys.org/news/2020-04-soundwaves-superfluid-parameter.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.