

Virus-infected honey bees more likely to gain entrance to healthy hives

April 27 2020, by Diana Yates



Entomology professor Adam Dolezal and his colleagues found that infection with the Israeli acute paralysis virus increases the likelihood that infected bees are accepted by foreign colonies. Credit: Fred Zwicky

Honey bees that guard hive entrances are twice as likely to allow in

trespassers from other hives if the intruders are infected with the Israeli acute paralysis virus, a deadly pathogen of bees, researchers report.

Their new study, reported in the *Proceedings of the National Academy of Sciences*, strongly suggests that IAPV [infection](#) alters honey bees' behavior and physiology in ways that boost the virus's ability to spread, the researchers say.

"The most important finding of our study is that IAPV infection increases the likelihood that infected bees are accepted by foreign colonies," said Adam Dolezal, a professor of entomology at the University of Illinois at Urbana-Champaign who led the new research. "Somehow, the infected bees are able to circumvent the guards of foreign colonies, which they shouldn't be able to do."

Previous studies have shown that IAPV-infected honey bees are more likely than healthy bees to lose their way when returning home from foraging trips. In commercial beekeeping operations where hives are stacked much closer together than in the wild, the virus is even more likely to spread from one infected colony to nearby healthy ones.

To capture the behavior of individual bees, researchers tagged each one with the equivalent of a QR code and continuously monitored their interactions. The scientists were able to simultaneously track the behaviors of as many as 900 bees.



Honey bees touch their mouthparts and antennae together to share food and information, but the practice also can transmit viruses. Credit: Fred Zwicky


In previous work, study co-author U. of I. entomologist Gene Robinson and his colleagues developed this automated system to study bees engaged in trophallaxis, a process by which honey bees exchange regurgitated food and other liquids. They used this system to study how IAPV infection might affect the bees' trophallaxis social network.

"Honey bees use trophallaxis to share food with each other as well as hormones and other signaling molecules that can affect their physiology and behavior. They do it in pairs by touching their mouthparts and antennae, and each bee does this with hundreds of partners a day," said Robinson, who directs the Carl R. Woese Institute for Genomic Biology

at Illinois. "Trophallaxis is essential to the spread of information and nutrition throughout the hive, but unfortunately, a behavior performed with such close social contact also allows [viral infections](#) to be transmitted through a hive."

In the new study, the scientists saw that honey bees altered their behavior in response to infection in their own hives. IAPV-infected bees—and bees that had had their immune systems stimulated to mimic infection—engaged in less trophallaxis than their healthy counterparts did.

The infected bees were just as mobile as the other bees, so their lower rates of trophallaxis were not the result of sluggishness from being sick, Dolezal said. The researchers believe this change in behavior is a general response to a health threat and not specific to IAPV infection, which is in line with previous research.



Researchers tagged each honey bee with the equivalent of a QR code and used

an automated system to study trophallaxis, a process by which the bees exchange regurgitated food and other liquids. The system allowed them to track how infection with IAPV affected the bees' trophallaxis social network. Credit: Tim Gernat

When the scientists placed honey bee workers at the entrance of a foreign hive, however, the infected bees engaged in more trophallaxis with the guards, the researchers found. The guards were more likely to admit them than to let in healthy bees or bees whose immune systems had been stimulated. This response was specific to IAPV infection.

"Something about them must be different," Dolezal said.

To test whether the IAPV-infected bees were giving off a different chemical odor than their healthy nest mates, the researchers analyzed the chemistry of the hydrocarbons that coat the bees' exoskeletons. They discovered distinct hydrocarbon profiles for healthy bees, IAPV-infected bees and immunostimulated bees.

"It seems that the virus is changing how the bees smell, and perhaps the [infected bees](#) also are behaving in a way that is meant to appease the guards by engaging more in trophallaxis," Dolezal said.

The new findings suggest that IAPV is evolving in ways that enhance its ability to infect as many hosts as possible, Dolezal said.

"If you're a virus, it's much more valuable to get transmitted to a new family group, like traveling from one city to a new city," he said. "And so how do you get there? You increase the chances that the sick bees leaving colony A are more likely to get into colony B."

More information: Amy C. Geffre et al., "Honey bee virus causes context-dependent changes in host social behavior," *PNAS* (2020).
www.pnas.org/cgi/doi/10.1073/pnas.2002268117

Provided by University of Illinois at Urbana-Champaign

Citation: Virus-infected honey bees more likely to gain entrance to healthy hives (2020, April 27)
retrieved 19 September 2024 from
<https://phys.org/news/2020-04-virus-infected-honey-bees-gain-entrance.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.