


# Strong X-ray pulsations detected from pulsar 3A 0726-260

May 5 2020, by Tomasz Nowakowski



The SXT lightcurve of 3A 0726-260 in 0.3-7 keV energy range. Credit: Roy et al., 2020.

Using AstroSat satellite, Indian astronomers have detected strong X-ray pulsations from an X-ray binary pulsar known as 3A 0726-260. The discovery, presented in a paper published April 26 on the arXiv pre-print repository, sheds more light on the nature of this poorly studied object.

X-ray pulsars (also known as accretion-powered pulsars) are sources displaying strict periodic variations in X-ray intensity, consisting of a magnetized neutron star in orbit with a normal stellar companion. In

these [binary systems](#), the X-ray emission is powered by the release of gravitational potential energy as material is accreted from a massive companion. X-ray pulsars are among the most luminous objects in the X-ray sky.

Located some 20,000 [light years](#) away, 3A 0726-260 (other designation 4U 0728-25) is one of the least-studied X-ray binary pulsars, despite being a persistent X-ray emitter at luminosity level of about 100 decillion ergs/s during its non-flaring states. The system has [orbital period](#) of approximately 34.55 days and contains an X-ray pulsar with spin period of about 103 seconds.

A team of astronomers led by Jayashree Roy of the University of Mumbai, India, investigated 3A 0726-260 in detail. Their study was based on the data from AstroSat's Large Area X-ray Proportional Counter (LAXPC) and Soft X-ray Telescope (SXT).

"We present results from broadband (0.3-40 keV) study of the source using data from AstroSat observations with the SXT and the LAXPC detectors," the astronomers wrote in the paper.

The observations detected strong X-ray pulsation with a period of 103.144 seconds. This suggests that there has been only a marginal change in the spin period of the [pulsar](#) since last measurements conducted in 1997. Moreover, weak pulsations from the source in 20-40 keV range were detected for the first time in the LAXPC data.

In general, the pulse profile of 3A 0726-260 was found to be energy dependent. It appears that the pulse shape changes from a broad single pulse, up to 5.0 keV, to a double pulse at higher energy (above 5.0 keV). The researchers offer two hypotheses that could explain such behavior.

"The change in the pulse profile from a single peak to a weak double

peaked structure may be explained by intrinsic change occurring in the beaming pattern from a pencil beam to a fan beam which results in the beam to move out of our line of sight (...). The change in the [pulse](#) profile can also be attributed to a transition in accretion pattern from a smooth accretion stream at low energies to several narrow accretion streams at high energy that are phase-locked with the neutron star," the paper reads.

The study also detected the presence of an of broad 1.06 keV, iron K-alpha line at approximately 6.3 keV. The astronomers added that 3A 0726-260 is accreting at almost a steady accretion rate with no indication of any instability that could trigger outbursts from this source.

**More information:** AstroSat observation of the Be/X-ray binary Pulsar 3A 0726-260 (4U 0728-25), arXiv:2004.12372 [astro-ph.HE] [arxiv.org/abs/2004.12372](https://arxiv.org/abs/2004.12372)

© 2020 Science X Network

Citation: Strong X-ray pulsations detected from pulsar 3A 0726-260 (2020, May 5) retrieved 22 September 2024 from <https://phys.org/news/2020-05-strong-x-ray-pulsations-pulsar-3a.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.