


Smaller farm fields can reduce biodiversity loss and increase wild plants, birds, beetles, and bats

July 10 2020, by Lenore Fahrig



Credit: AI-generated image ([disclaimer](#))

Around the world, biodiversity is dropping precipitously. In North America, about [three billion birds](#) have been lost over recent decades and [amphibian populations](#) are declining at a rate of about four percent per year. Globally, [insects are vanishing](#) at a rate of about nine percent

per decade.

This is distressing for nature lovers. It is also worrying on a practical level: in agricultural landscapes, [wild species](#) perform many important activities such as pollination and control of insect pests.

A highly effective way to reduce [biodiversity](#) losses in agricultural regions is to increase the amount of natural and semi-natural habitats such as forests, hedgerows, prairie strips and other non-crop areas. This has led to various government [incentives for farmers to leave some areas out of production](#). But [there are limits](#) to how much land farmers can leave for biodiversity, while foregoing opportunities for [crop production](#).

The benefit of small crop fields

Small fields can help. [Farmers can maintain or increase biodiversity](#) in agricultural landscapes by [planting their crops in smaller fields](#).

For example, instead of planting two large fields, they could plant eight small fields, keeping the same total amount of land in production. Note that this is not about the size of a farm, but about the average size of crop fields.

A very large farm could, in theory, be composed of a very large number of small fields, where a "field" is an area in a particular crop type, such as a corn field or a hay field. Adjacent areas in the same crop type are separate fields if they are divided by even a narrow bit of non-field such as a hedgerow, a fence line, a track or a ditch.



Credit: AI-generated image ([disclaimer](#))

My colleagues and I have found remarkably consistent increases in biodiversity with decreasing average field size, across eight very different agricultural regions in five countries—and for a wide array of wildlife types.

The number of [species of plants](#), [pollinators including bees, butterflies and syrphid flies](#), and pest-eating wildlife such as spiders, carabid beetles, birds, [frogs](#) and [bats](#), all increase with decreasing average field size. For example, biodiversity is about 50 percent higher when the average field size drops to two hectares from eight hectares.

In fact, the benefit to biodiversity of reducing crop field sizes appears to be greater than two other commonly suggested methods for increasing biodiversity in agricultural landscapes—[reducing pesticide use](#) and

increasing crop diversity, the number of different crop types grown in a landscape.

The best of both worlds

One question is whether the increased biodiversity is due to the additional non-cropped strips of semi-natural land, such as hedgerows, between fields. In other words, is there actually less cropland—and more semi-natural land—in an agricultural landscape with eight small fields than in a landscape with two large ones?

Our research shows that in fact biodiversity is higher in a [landscape](#) with smaller crop fields, even for the same total amount of natural and semi-natural habitat, including all the little pieces such as hedgerows.

We don't know yet why biodiversity is higher in landscapes with smaller crop fields. One possibility is that when crop fields are smaller, the natural habitats are closer together.



Credit: AI-generated image ([disclaimer](#))

This would increase the ability of wildlife to [easily access](#) natural habitats from crop fields. Crop fields may provide temporary food sources for wildlife, such as when a crop is in flower or when there is a pest outbreak. Natural areas are also needed for breeding and overwintering sites and for escape from farming operations such as plowing.

When wildlife can easily move back and forth between [crops](#) and natural spaces, perhaps they have access to the best of both worlds.

The farmer's dilemma

Altogether, [our research suggests](#) a "[many small](#)" approach to support

biodiversity in [agricultural landscapes](#).

But can farmers readily reduce the sizes of fields? In many [parts of the world, just the opposite](#) has been happening over the past 50 years.

Fields are being consolidated by removing the tracks, fences, stone walls and hedgerows that once separated them. Farmers do this to increase the efficiency of their operations. But in doing so, they reduce the populations of wildlife that help support agriculture.

The net cost or benefit to farmers of reducing field sizes while maintaining the same area in production has not yet been studied. But the benefits to biodiversity are now clear.

This article is republished from [The Conversation](#) under a Creative Commons license. Read the [original article](#).

Provided by The Conversation

Citation: Smaller farm fields can reduce biodiversity loss and increase wild plants, birds, beetles, and bats (2020, July 10) retrieved 22 September 2024 from <https://phys.org/news/2020-07-smaller-farm-fields-biodiversity-loss.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.