


# Researchers identify evolutionary origins of SARS-CoV-2

July 28 2020

---



Credit: Pixabay/CC0 Public Domain

By reconstructing the evolutionary history of SARS-CoV-2, the virus that is responsible for the COVID-19 pandemic, an international research team of Chinese, European, and U.S. scientists has discovered

that the lineage that gave rise to the virus has been circulating in bats for decades and likely includes other viruses with the ability to infect humans. The findings, which University of Glasgow scientists contributed to, have implications for the prevention of future pandemics stemming from this lineage.

"Coronaviruses have [genetic material](#) that is highly recombinant, meaning different regions of the [virus's](#) genome can be derived from multiple sources," said Maciej Boni, associate professor of biology, Penn State. "This has made it difficult to reconstruct SARS-CoV-2's origins. You have to identify all the regions that have been recombining and trace their histories. To do that, we put together a diverse team with expertise in recombination, phylogenetic dating, virus sampling, and molecular and viral evolution."

The team used three different bioinformatic approaches to identify and remove the recombinant regions within the SARS-CoV-2 genome. Next, they reconstructed phylogenetic histories for the non-recombinant regions and compared them to each other to see which specific viruses have been involved in recombination events in the past. They were able to reconstruct the evolutionary relationships between SARS-CoV-2 and its closest known bat and pangolin viruses. Their findings appear today in *Nature Microbiology*.

The researchers found that the lineage of viruses to which SARS-CoV-2 belongs diverged from other bat viruses about 40-70 years ago. Importantly, although SARS-CoV-2 is genetically similar (about 96%) to the RaTG13 coronavirus, which was sampled from a *Rhinolophus affinis* horseshoe bat in 2013 in Yunnan province, China, the team found that it diverged from RaTG13 a relatively long time ago, in 1969.

"The ability to estimate divergence times after disentangling recombination histories, which is something we developed in this

collaboration, may lead to insights into the origins of many different viral pathogens," said Philippe Lemey, principal investigator in the Department of Evolutionary and Computational Virology, KE Leuven.

The team found that one of the older traits that SARS-CoV-2 shares with its relatives is the receptor-binding domain (RBD) located on the Spike protein, which enables the virus to recognize and bind to receptors on the surfaces of human cells.

"This means that other viruses that are capable of infecting humans are circulating in [horseshoe bats](#) in China," said David L. Robertson, professor of computational virology, MRC-University of Glasgow Center for Virus Research.

Will these viruses be capable of jumping directly from bats into humans or will an intermediate species be required to make the leap? According to Robertson, for SARS-CoV-2, other research groups incorrectly proposed that key evolutionary changes occurred in pangolins.

"SARS-CoV-2's RBD sequence has so far only been found in a few pangolin viruses," said Robertson. "Furthermore, the other key feature thought to be instrumental to SARS-CoV-2's ability to infect humans—a polybasic cleavage site insertion in the Spike protein—has not yet been seen in another close bat relative of the SARS-CoV-2 virus. Yet, while it is possible that pangolins may have acted as an intermediate host facilitating transmission of SARS-CoV-2 to humans, no evidence exists to suggest that pangolin infection is a requirement for bat viruses to cross into humans. Instead, our research suggests that SARS-CoV-2 likely evolved the ability to replicate in the upper respiratory tract of both humans and pangolins."

The team concluded that preventing future pandemics will require better sampling within wild bats and the implementation of human disease

surveillance systems that are able to identify novel pathogens in humans and respond in real time.

"The key to successful surveillance," said Robertson, "is knowing which viruses to look for and prioritizing those that can readily infect humans. We should have been better prepared for a second SARS virus."

Boni added, "We were too late in responding to the initial SARS-CoV-2 outbreak, but this will not be our last coronavirus pandemic. A much more comprehensive and real-time surveillance system needs to be put in place to catch viruses like this when case numbers are still in the double digits."

**More information:** Maciej F. Boni et al. Evolutionary origins of the SARS-CoV-2 sarbecovirus lineage responsible for the COVID-19 pandemic, *Nature Microbiology* (2020). [DOI: 10.1038/s41564-020-0771-4](https://doi.org/10.1038/s41564-020-0771-4)

Provided by University of Glasgow

Citation: Researchers identify evolutionary origins of SARS-CoV-2 (2020, July 28) retrieved 21 September 2024 from <https://phys.org/news/2020-07-evolutionary-sars-cov-.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.