


# Cephalopod *Sinoceras chinense* discovered outside China for the first time

30 July 2020, by Li Yuan



large number of fossil specimens.

They discovered *S. chinense* in the Upper Ordovician strata, which is the first discovery outside China. The study was published in *Palaeoworld* on June 18.

These specimens were recorded in a geoconservation site named Nautiloid Site, located in western Thailand. Previously, these specimens were wrongly identified as actinocerids, leading to miscalculation of the geological time of the upper part of Tha Manao Formation.


The exact identification of *S. chinense* suggested that the age of the upper part of Tha Manao Formation was the early Katian of Late Ordovician, providing evidence for the stratigraphic correlation between the upper part of the Tha Manao Formation in Thailand and the Pagoda Formation (and contemporaneous units) in China.

*Sinoceras chinense*, collected from Si Sawat county of Kanchanabura Province, western Thailand. Credit: NIGPAS

*Sinoceras chinense* is a kind of extinct cephalopod mollusk that flourished in the late Ordovician. It is regarded as the index fossil of the Upper Ordovician Pagoda Formation on the Yangtze Platform of South China, with a likely age of early Katian.

*S. chinense* was previously only known from the Upper Ordovician of Chinese blocks/terranes, including South China, Tarim, Tibet and western Yunnan.

In January 2020, a Sino-Thai research team led by Dr. Fang Xiang from the Nanjing Institute of Geology and Palaeontology of the Chinese Academy of Sciences (NIGPAS) and Prof. Clive Burrett from Mahasarakham University conducted [field research](#) in western Thailand and collected a



Distribution of *S. chinense* in the northeastern peri-Gondwana region. Credit: NIGPAS

Western Thailand, combined with Baoshan region of western Yunnan, was located on Sibumasu Terrane during the Early Palaeozoic. The confirmed identification of *S. chinense* in Thailand is the first record in Thailand and also the first report in a region outside of China.

Moreover, the discovery of the species in the Sibumasu Terrane provides strong support for the palaeogeographic reconstruction and pronounced palaeobiogeographic changes from the Middle to Late Ordovician among the peri-Gondwanan regions.

**More information:** Xiang Fang et al. *Sinoceras chinense* (Foord, 1888) in western Thailand: first identification outside China, *Palaeoworld* (2020).  
[DOI: 10.1016/j.palwor.2020.06.004](https://doi.org/10.1016/j.palwor.2020.06.004)

Provided by Chinese Academy of Sciences

APA citation: Cephalopod *Sinoceras chinense* discovered outside China for the first time (2020, July 30) retrieved 20 September 2020 from <https://phys.org/news/2020-07-cephalopod-sinoceras-chinense-china.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*