


Topological superconducting phase protected by 1-D local magnetic symmetries

27 August 2020



Credit: CC0 Public Domain

Topological superconductors (TSCs) are new kind of topological quantum states with fully superconducting gapped band structure in the bulk, but they support gapless excitations called Majorana zero modes (MZMs) at the boundaries. Because of their nonlocal correlation and non-Abelian statistic nature, MZMs are proposed as the qubits of topological quantum computation. Hence, searching and operating the MZMs in TSC materials is now an important topic in condensed matter physics.

To identify a TSC, one should first ascertain its topological classification. The topological classification highly depends on the symmetries including [time-reversal symmetry](#), particle-hole symmetry, and especially the crystalline symmetries. Without the consideration of crystalline symmetries, the Bogoliubov-deGennes (BdG) Hamiltonians of the 1-D superconductors only have the Z2 classification. The mirror reflection symmetry and rotational symmetries can enhance the classification to Z class. Nevertheless, the topological classification of superconductors with general magnetic symmetries is still an open

question.

In a new research article published in the Beijing-based *National Science Review*, scientists from the Huazhong University of Science and Technology in Wuhan, China, and Princeton University in New Jersey, USA proposed the method to classify the topological superconducting phase by examining the compatibility between different MZMs. Co-authors Jinyu Zou, Qing Xie, Zhida Song and Gang Xu analyzed the topological classification of gapped superconducting wires with local magnetic symmetries (LMSs). They found that an effective BDI class TSC can be realized in M_xT or $C_{2z}T$ invariant wire. Remarkably, the new TSC phases characterized by Z^h invariant in $C_{4z}T$ case and $Z^h \oplus Z^c$ invariant in $C_{6z}T$ case are discovered.

T'	M_xT ($n = 2$)	$C_{2z}T$ ($n = 2$)	$C_{4z}T$ ($n = 4$)	$C_{6z}T$ ($n = 6$)
T^n	1	1	-1	1
P^2	1	1	1	1
S^n	1	1	-1	1
Invariant	Z^c (BDI)	Z^c (BDI)	Z^h ($2 \times AIII$)	$Z^h \oplus Z^c$ ($2 \times AIII \oplus BDI$)

The topological classification of the 1D gapped superconducting systems with the LMSs. Credit: ©Science China Press


In the article titled "New types of [topological superconductors](#) under local magnetic symmetries". The authors focus on the 1D superconducting wires with LMSs $T' = M_xT, C_{2z}T, C_{4z}T$ and $C_{6z}T$. "The operation of T' does not change the position of electrons. Hence it acts on the BdG Hamiltonian like a time-reversal operator". Combining T' and particle-hole symmetry P leads to a chiral symmetry $S = T'P$. The BdG Hamiltonian can adopt the

diagonalized form according to the chiral symmetry. And the MZMs are the eigenstates of the chiral symmetry S . The authors find "MZMs having chiral eigenvalues s and $-s$ can couple to each other and be eliminated." Following the guideline, they analysis the compatibility of the MZMs at the end of the 1D superconducting wires with LMSs, and summarize their topological classification as listed in Table I.

one can couple the 1D TSCs in y direction to construct a 2-D TSC. The high symmetry lines $k_y = 0$ and $k_y = \pi$ in momentum space preserve the 1D LMS. With proper parameters, the $k_y = 0$ and $k_y = \pi$ lines can belong to distinct topological phase, and result in the gapless propagating Majorana edge states connecting the conducting bands and valence bands."

More information: Jinyu Zou et al, New types of topological superconductors under local magnetic symmetries, *National Science Review* (2020). DOI: [10.1093/nsr/nwaa169](https://doi.org/10.1093/nsr/nwaa169)

Provided by Science China Press



(a) A C4zT-preserving superconducting wire aligned along z direction. (b) The topological phase diagram. (c) the spectrum of nontrivial phase with an open boundary on both sides, in which four MZMs appear at zero energy. Credit: ©Science China Press

"To illustrate the TSC phase with the LMS $C_{4z}T$, we construct a 1D anti-ferromagnetic chain along z-direction," the scientists add. They give the topological phase diagram of the model. "In the nontrivial TSC phase, the open quantum wire traps an integer pairs of MZMs at its ends." they also show the MZMs through numerical and analytical calculation.

"These results not only enrich the variety of the 1-D TSC, but also provide luxuriant building blocks for the construction of new type 2-D and 3-D TSCs" they forecast at the end of the article, "For example,

APA citation: Topological superconducting phase protected by 1-D local magnetic symmetries (2020, August 27) retrieved 17 January 2021 from <https://phys.org/news/2020-08-topological-superconducting-phase-d-local.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.