


# High-fidelity record of Earth's climate history puts current changes in context

September 10 2020



Past and future trends in global mean temperature spanning the last 67 million years. Oxygen isotope values in deep-sea benthic foraminifera from sediment cores are a measure of global temperature and ice volume. Temperature is relative to the 1961-1990 global mean. Data from ice core records of the last 25,000 years illustrate the transition from the last glacial to the current warmer period, the Holocene. Historic data from 1850 to today show the distinct increase after 1950 marking the onset of the Anthropocene. Future projections for global temperature for three Representative Concentration Pathways (RCP) scenarios in relation to the benthic deep-sea record suggest that by 2100 the climate state will be comparable to the Miocene Climate Optimum (~16 million years ago), well beyond the threshold for nucleating continental ice sheets. If emissions are constant after 2100 and are not stabilized before 2250, global climate by 2300 might enter the hothouse world of the early Eocene (~50 million years ago) with its multiple global warming events and no large ice sheets at the

poles. Credit: Westerhold et al., CENOGRID

For the first time, climate scientists have compiled a continuous, high-fidelity record of variations in Earth's climate extending 66 million years into the past. The record reveals four distinctive climate states, which the researchers dubbed Hothouse, Warmhouse, Coolhouse, and Icehouse.

These major [climate](#) states persisted for millions and sometimes tens of millions of years, and within each one the climate shows rhythmic variations corresponding to changes in Earth's orbit around the sun. But each climate state has a distinctive response to orbital variations, which drive relatively small changes in [global temperatures](#) compared with the dramatic shifts between different climate states.

The new findings, published September 10 in *Science*, are the result of decades of work and a large international collaboration. The challenge was to determine past climate variations on a [time scale](#) fine enough to see the variability attributable to orbital variations (in the eccentricity of Earth's orbit around the sun and the precession and tilt of its rotational axis).


"We've known for a long time that the glacial-interglacial cycles are paced by changes in Earth's orbit, which alter the amount of solar energy reaching Earth's surface, and astronomers have been computing these orbital variations back in time," explained coauthor James Zachos, distinguished professor of Earth and planetary sciences and Ida Benson Lynn Professor of Ocean Health at UC Santa Cruz.

"As we reconstructed past climates, we could see long-term course changes quite well. We also knew there should be finer-scale rhythmic variability due to orbital variations, but for a long time it was considered

impossible to recover that signal," Zachos said. "Now that we have succeeded in capturing the natural climate variability, we can see that the projected anthropogenic warming will be much greater than that."

For the past 3 million years, Earth's climate has been in an Icehouse state characterized by alternating glacial and interglacial periods. Modern humans evolved during this time, but [greenhouse gas emissions](#) and other human activities are now driving the planet toward the Warmhouse and Hothouse climate states not seen since the Eocene epoch, which ended about 34 million years ago. During the early Eocene, there were no polar ice caps, and average global temperatures were 9 to 14 degrees Celsius higher than today.

"The IPCC projections for 2300 in the 'business-as-usual' scenario will potentially bring global temperature to a level the planet has not seen in 50 million years," Zachos said.



The new global climate record CENOGRID (lower panel) is the first to continually and accurately trace how Earth's climate has changed since the great extinction of the dinosaurs 66 million years ago. The record was generated using the oxygen (shown) and carbon isotopes from tiny microfossils found in deep-sea sediments collected by the IODP ship R/V JOIDES Resolution (shown in the photo) and shows the natural range of climate change and variability over the last 66 million years. Credit: Thomas Westerhold / Adam Kutz

Critical to compiling the new climate record was getting high-quality sediment cores from deep ocean basins through the international Ocean Drilling Program (ODP, later the Integrated Ocean Drilling Program, IODP, succeeded in 2013 by the International Ocean Discovery Program). Signatures of past climates are recorded in the shells of microscopic plankton (called foraminifera) preserved in the seafloor sediments. After analyzing the sediment cores, researchers then had to develop an "astrochronology" by matching the climate variations recorded in sediment layers with variations in Earth's orbit (known as Milankovitch cycles).

"The community figured out how to extend this strategy to older time intervals in the mid-1990s," said Zachos, who led a study published in 2001 in *Science* that showed the climate response to orbital variations for a 5-million-year period covering the transition from the Oligocene epoch to the Miocene, about 25 million years ago.

"That changed everything, because if we could do that, we knew we could go all the way back to maybe 66 million years ago and put these transient events and major transitions in Earth's climate in the context of orbital-scale variations," he said.

Zachos has collaborated for years with lead author Thomas Westerhold at the University of Bremen Center for Marine Environmental Sciences (MARUM) in Germany, which houses a vast repository of sediment cores. The Bremen lab along with Zachos's group at UCSC generated much of the new data for the older part of the record.

Westerhold oversaw a critical step, splicing together overlapping segments of the climate record obtained from sediment cores from different parts of the world. "It's a tedious process to assemble this long megasplice of climate records, and we also wanted to replicate the records with separate sediment cores to verify the signals, so this was a

big effort of the international community working together," Zachos said.

Now that they have compiled a continuous, astronomically dated climate record of the past 66 million years, the researchers can see that the climate's response to orbital variations depends on factors such as greenhouse gas levels and the extent of polar ice sheets.

"In an extreme greenhouse world with no ice, there won't be any feedbacks involving the ice sheets, and that changes the dynamics of the climate," Zachos explained.

Most of the major climate transitions in the past 66 million years have been associated with changes in greenhouse gas levels. Zachos has done extensive research on the Paleocene-Eocene Thermal Maximum (PETM), for example, showing that this episode of rapid global warming, which drove the climate into a Hothouse state, was associated with a massive release of carbon into the atmosphere. Similarly, in the late Eocene, as atmospheric carbon dioxide levels were dropping, ice sheets began to form in Antarctica and the climate transitioned to a Coolhouse state.

"The climate can become unstable when it's nearing one of these transitions, and we see less predictable responses to orbital forcing, so that's something we would like to better understand," Zachos said.

The new climate record provides a valuable framework for many areas of research, he added. It is not only useful for testing climate models, but also for geophysicists studying different aspects of Earth dynamics and paleontologists studying how changing environments drive the evolution of species.

"It's a significant advance in Earth science, and a major legacy of the

international Ocean Drilling Program," Zachos said.

**More information:** T. Westerhold et al., "An astronomically dated record of Earth's climate and its predictability over the last 66 million years," *Science* (2020). [science.sciencemag.org/cgi/doi ... 1126/science.aba6853](https://science.sciencemag.org/cgi/doi/10.1126/science.aba6853)

Provided by University of California - Santa Cruz

Citation: High-fidelity record of Earth's climate history puts current changes in context (2020, September 10) retrieved 20 September 2024 from <https://phys.org/news/2020-09-high-fidelity-earth-climate-history-current.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--