

Animals lose fear of predators rapidly after they start encountering humans

September 22 2020



Credit: Unsplash/CC0 Public Domain

Most wild animals show a suite of predator avoidance behaviors such as vigilance, freezing, and fleeing. But these are quickly reduced after the animals come into contact with humans through captivity, domestication, or urbanization, according to a study led by Benjamin Geffroy from MARBEC (Institute of Marine Biodiversity, Exploitation and Conservation), publishing September 22nd in the open-access journal

PLOS Biology.

The international team of researchers analyzed the results of 173 peer-reviewed studies investigating antipredator traits (behavioral and physiological) in 102 species of domesticated, captive, and urbanized mammals, birds, reptiles, fish and molluscs, while taking into account their position in the Tree of Life.

The scientists found that contact with humans led to a rapid loss of animals antipredator traits, but simultaneously the variability between individuals initially increases and then gradually decreases over the generations in contact with [human](#). The authors suppose that this two-step process is caused by reduced pressure from [natural selection](#) as a result of living in a safer environment, followed by [artificial selection](#) by humans for docility in the case of domestication.

Animals showed immediate changes in antipredator responses in the first generation after contact with humans, suggesting that the initial response is a result of behavioral flexibility, which may later be accompanied by genetic changes if contact continues over many generations. The researchers also found that domestication altered animal antipredator responses three times faster than urbanization, while captivity resulted in the slowest changes. The results also showed that herbivores changed behavior more rapidly than carnivores and that solitary species tended to change quicker than group-living animals.

The study demonstrates that domestication and urbanization exert similar pressures on animals and can result in rapid behavioral changes. The loss of anti-predator behaviors can cause problems when those domesticated or urbanized species encounter predators or when [captive animals](#) are released back into the wild. Understanding how animals respond to contact with humans has important implications for conservation and [urban planning](#), captive breed programs, and livestock management.

Dr. Geffroy adds "While it is well known that the fact of being protected by humans decreases antipredator capacities in [animals](#), we did not know how fast this occurs and to what extent this is comparable between contexts! We also integrated physiological traits in the study but they were much less numerous than behavioral traits. We believe they should be systematically investigated to draw a global pattern of what is happening at the individual level. We need more data to understand whether this occurs also with the mere presence of tourists."

More information: Geffroy B, Sadoul B, Putman BJ, Berger-Tal O, Garamszegi LZ, Møller AP, et al. (2020) Evolutionary dynamics in the Anthropocene: Life history and intensity of human contact shape antipredator responses. *PLoS Biol* 18(9): e3000818.
doi.org/10.1371/journal.pbio.3000818

Provided by Public Library of Science

Citation: Animals lose fear of predators rapidly after they start encountering humans (2020, September 22) retrieved 18 April 2024 from <https://phys.org/news/2020-09-animals-predators-rapidly-encountering-humans.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.