


# Rubber-leguminous shrub systems should be popularized to reduce greenhouse gas emissions in Xishuangbanna

October 14 2020, by Zhang Nannan

---



Rubber-*Flemingia macrophylla* plantation in Xishuangbanna. Credit: LIU Changan

In recent years, numerous rubber-based agroforestry systems, using a biological approach to enhance ecosystem services, have been developed. Intercropping rubber with *Flemingia macrophylla*, a

leguminous shrub, has been widely established in rubber plantations area of China.

Previous studies showed that [greenhouse gas emissions](#) in [rubber](#) and rubber-F. macrophylla systems would be affected by their [plantation](#) age. However, the effect of different-aged rubber and rubber-F. macrophylla systems on CO<sub>2</sub>, N<sub>2</sub>O, and CH<sub>4</sub> emissions from soils is unknown.

In a study published in *Forest Ecology and Management*, researchers from the Xishuangbanna Tropical Botanical Garden (XTBG) investigated the effect of introducing F. macrophylla to different-aged [rubber plantations](#) on the emissions of CO<sub>2</sub>, N<sub>2</sub>O, and CH<sub>4</sub> in Xishuangbanna.

They found that the emissions of CO<sub>2</sub>, N<sub>2</sub>O and CH<sub>4</sub> decreased as the trees aged in the rubber plantations. The introduction of F. macrophylla to different-aged rubber plantations significantly decreased CO<sub>2</sub> and CH<sub>4</sub> flux but increased N<sub>2</sub>O flux.

Furthermore, the CO<sub>2</sub> and N<sub>2</sub>O fluxes were mainly affected by [soil](#) temperature at 10 cm depth, and CH<sub>4</sub> flux was mainly affected by both soil water content in the 0–10 cm soil layer and soil temperature at 10 cm depth.

Moreover, they found that the rubber-leguminous shrub systems significantly improved soil organic carbon sequestration rate and reduced greenhouse gas emissions, relative to the same-aged rubber plantations.



Sloping fields in the rubber–*Flemingia macrophylla* plantations. Credit: LIU Changan

Decrease of CO<sub>2</sub> emissions is crucial for the reduction of greenhouse gas emissions in rubber plantations worldwide.

"We therefore propose that rubber-leguminous shrub systems should be popularized in the rubber planting area for reducing greenhouse gas emissions," said Prof. Tang Jianwei, principal investigator of the study.

**More information:** Xin Rao et al. Rubber-leguminous shrub systems stimulate soil N<sub>2</sub>O but reduce CO<sub>2</sub> and CH<sub>4</sub> emissions, *Forest Ecology and Management* (2020). [DOI: 10.1016/j.foreco.2020.118665](https://doi.org/10.1016/j.foreco.2020.118665)

Provided by Chinese Academy of Sciences

Citation: Rubber-leguminous shrub systems should be popularized to reduce greenhouse gas emissions in Xishuangbanna (2020, October 14) retrieved 19 April 2024 from <https://phys.org/news/2020-10-rubber-leguminous-shrub-popularized-greenhouse-gas.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.