


# Seesaw of Indo-Pacific summer monsoons triggered by the tropical Atlantic Ocean

29 October 2020



The increasing influences from the tropical Atlantic sea surface temperature could trigger the observed multidecadal seesaw of Indo-Pacific summer monsoons in terms of their intensity of interannual variability and monsoon-ENSO biennial relationship variability. Credit: Dr. Lei WANG

The Indian summer monsoon (ISM) and western North Pacific summer monsoon (WNPSM) are two major subcomponents of the Asian summer monsoon. The monsoon rainfall is the life-blood of more than half the world's population. A better understanding of the ISM and WNPSM variability is of vital importance both in a socio-economic and scientific sense.

In a recent study published in *Atmospheric and Oceanic Science Letters*, the authors, from Guangdong Ocean University, quantitatively examined and compared the intensity of the interannual variability (IIV) of the monsoon and monsoon-ENSO biennial relationship (MEBR) for both the ISM and WNPSM.

"Our results reveal interesting multidecadal seesaw patterns of the ISM and WNPSM in terms of their IIV and MEBR, which could be triggered by the increasing influences from the tropical Atlantic Ocean," says Dr. WANG Lei, the corresponding author of this study.

Out-of-phase covariability was observed between

the ISM and WNPSM. When the IIV and MEBR of the ISM were strong (weak) before (after) the mid-1980s, the IIV and MEBR of the WNPSM tended to be weak (strong). During the period with a stronger ENSO-Atlantic (ENSO-Indian Ocean) coupling, the IIV and MEBR of the WNPSM (ISM) were observed to be stronger. The increasing influences from the tropical Atlantic Ocean sea surface temperature (SST) could exert different influences on the ISM and WNPSM, leading to changes in opposite signs of the two monsoons in terms of IIV and MEBR. Increasing influences from the tropical Atlantic SST may be related to its persistent warming trend due to combined effects from the global warming trend and the recent phase change of the Atlantic multidecadal oscillation.

"These results imply that tropical Atlantic SST may need to be given more attention when predicting future [monsoon](#) variability of the ISM and WNPSM," concludes Dr. WANG. "More studies are needed in the future to better understand the inter-basin interactions among the tropical Pacific, Atlantic, and Indian Ocean basins, and their interdecadal changes, which may be the key to understanding the recent climate variability."

**More information:** Baiyang CHEN et al, Contrasting the Indian and western North Pacific summer monsoons in terms of their intensity of interannual variability and biennial relationship with ENSO, *Atmospheric and Oceanic Science Letters* (2020). [DOI: 10.1080/16742834.2020.1806683](https://doi.org/10.1080/16742834.2020.1806683)

Provided by Chinese Academy of Sciences

APA citation: Seesaw of Indo-Pacific summer monsoons triggered by the tropical Atlantic Ocean (2020, October 29) retrieved 27 January 2021 from <https://phys.org/news/2020-10-seesaw-indo-pacific-summer-monsoons-triggered.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*