


Uncovering the hidden side of storms: France's Taranis satellite to launch in November

November 19 2020, by Manon Colonna D'istria



Credit: CNRS

Sprites, elves, jets... few people know that scientists habitually use such other-worldly words to describe transient luminous events or TLEs, light flashes that occur during active storms just a few tens of kilometers over

our heads. Few people also know that storms can act as particle accelerators generating very brief bursts of X-rays and gamma rays. But what are the physical processes and mechanisms behind these phenomena discovered barely 30 years ago? Do they impact the physics and chemistry of the upper atmosphere, the environment or even humans? Such are the questions facing the French Taranis satellite that will be riding aloft during the night of 16 to 17 November atop a Vega launcher from the Guiana Space Center, an all-French mission involving research scientists from CNES, the national scientific research center CNRS, the atomic energy and alternative energies commission CEA and several French universities.

TLEs and terrestrial gamma-ray flashes (TGFs) are seen all over the world where storms occur. But because we don't know enough about them, they don't feature in the toolbox of climatologists and meteorologists. Are they implicated in the increasing number of extreme weather events? If so, they could be modeled and factored into forecasts in real time. Although Taranis is first and foremost a fundamental research satellite, the data it is set to deliver on Earth's thermal and climate mechanisms could serve more operational applications like climatology and weather forecasting.

Elves, sprites, sprite halos, blue jets and even pixies or gnomes are just some of the whimsical names given to the range of phenomena in the generic family of TLEs—a poetic lexicon that contrasts sharply with their violence. These ephemeral upper-atmosphere events occur between the tops of storm clouds and an altitude of 90 kilometers. First predicted as early as 1920, their existence was not confirmed until the nineties. They have since been recorded by numerous ground and space observations. Elves take the form of an expanding glow of light, appearing at an altitude of 90 kilometers and lasting no more than one millisecond; an active storm may produce thousands of them in the space of a few hours. Occurring between 40 and 90 kilometers above Earth's

surface, sprites have a complex structure of branches and tendrils and can last for up to 10 milliseconds. Blue jets appear at the top of storm clouds and propagate to altitudes of up to 50 kilometers. Occasionally, 'gigantic' jets may propagate up to 90 kilometers.

TGFs were first observed scientifically in 1994 by the Compton Gamma-Ray Observatory (CGRO), a NASA spacecraft deployed from the U.S. space shuttle Atlantis. In certain conditions, storms generate a very short burst of gamma photons. TGFs were for a time considered a rare occurrence accompanying sprites; we now know they are generated by electric activity in clouds. For lack of the right instruments, the Italian AGILE satellite (2007) and U.S. Fermi space telescope (2008) were unable to fully confirm current hypotheses on the mechanisms that generate them or estimate their number. Taranis will therefore bring new insights into how they are generated and their radiation impact, which has never been measured before.

In France, the [atomic energy](#) agency CEA first turned its attention to these transient events and their impact in 1993. On 9 December 2010, the project got the official go-ahead from CNES's Board of Directors. Taranis is an all-French mission with science goals set by French research laboratories. In addition to CEA, CNRS is closely involved through several of its affiliated research laboratories¹: the LPC2E environmental and space physics and chemistry laboratory is coordinating development of the science payload, is responsible for the science mission center and is contributing instruments; the IRAP astrophysics and planetology research institute, the LATMOS atmospheres, environments and space observations laboratory and the APC astroparticles and cosmology laboratory are contributing to the payload.




Illustration of the Taranis. Credit: CNES/ill./SATTLER Oliver, 2012

Other instruments on Taranis include outside contributions from Stanford University and Goddard Space Flight Center (GSFC) in the United States, the Institute of Atmospheric Physics (IAP) and Charles University in the Czech Republic and the Space Research Center of the Polish Academy of Sciences (CBK).

Taranis looks somewhat different, as in place of the aluminized or gold-plated Mylar insulation traditionally used on satellites it is coated with a special black and white paint. This is not just attention to esthetic detail, the purpose of the paint being to avoid interfering with the surrounding [electric field](#) and prevent reflected light disrupting the optical sensors. A less visible but key feature is the original design of its payload, comprising eight instruments operated as a single unit thanks to MEXIC, the brain of Taranis that powers and synchronizes the instruments and manages the payload, executes the trigger strategy to capture an event and even handles the transfer of selected data to mass memory.

Taranis' payload close up:

- XGRE: three X-ray and gamma-ray detectors for measuring high-energy photons (50 keV to 10 MeV) and relativistic electrons (1 MeV to 10 MeV) – APC/IRAP/CNES
- MCP (MC-U and PH-U): two cameras (10 images per second) and four photometers to measure luminance in different spectral bands—CEA/CNES
- IDEE: two high-energy electron detectors (70 keV to 4 MeV) – IRAP/Charles University
- IMM: three-axis magnetometer to measure the alternating magnetic field (5 Hz to 1 MHz) – LPC2E/Stanford University
- IME-HF: HF antenna for measuring the high-frequency electric field (100 kHz to 35 MHz) – LPC2E/IAP
- IME-BF: instrument for measuring the low-frequency electric field (DC to 1 MHz) – LATMOS
- SI: ion probe to determine thermal plasma fluctuations—GSFC/LATMOS
- MEXIC: two electronic units comprising eight analysers, each connected to an instrument. It powers each instrument, handles payload modes and interfaces with mass memory and the onboard computer. MEXIC will also be tasked with synchronizing the instruments when events are detected (TLEs by MCP's photometers, TGFs by XGRE, electron beams by IDEE, wave bursts by IME-HF) – LPC2E/CBK

For two to four years, Taranis will scan regions of the sky where storm activity is intense and the probability of seeing TLEs and/or TGFs high. While it may be a national program, its results are eagerly awaited by the wider international scientific community. In atmospheric chemistry and physics, environmental science, climatology, high-energy astrophysics and many more fields besides, Taranis is set to reveal new insights—and science efforts won't end there, as the mission will undoubtedly pave the

way for future investigations.

Provided by CNRS

Citation: Uncovering the hidden side of storms: France's Taranis satellite to launch in November (2020, November 19) retrieved 21 September 2024 from
<https://phys.org/news/2020-11-uncovering-hidden-side-storms-france.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.