


Confirming simulated calculations with experiment results

19 November 2020



Credit: The University of Hong Kong

Dr. Zi Yang Meng from the Division of Physics and Astronomy, Faculty of Science, the University of Hong Kong (HKU), is pursuing a new paradigm of quantum material research that combines theory, computation and experiment in a coherent manner. Recently, he teamed up with Dr. Wei LI from Beihang University, Professor Yang Qi from Fudan University, Professor Weiqiang YU from Renmin University and Professor Jinsheng Wen from Nanjing University to untangle the puzzle of Nobel Prize-winning theory Kosterlitz-Thouless (KT) phase.

Not long ago, Dr. Meng, Dr. Li and Dr. Qi achieved accurate model calculations of a topological KT phase for a rare-earth magnet TmMgGaO_4 (TMGO), by performing computation on the Supercomputers Tianhe 1 and Tianhe 2; this time, the team overcame several conceptual and experimental difficulties, and succeeded in discovering a topological KT phase and its transitions in the same rare-earth magnet via highly sensitive nuclear magnetic resonance (NMR) and magnetic susceptibility measurements, means of detecting magnetic responses of material. The former one is more sensitive in detecting small magnetic moments while the latter one can facilitate easy implementation of the experiment.

These [experimental results](#), further explained the

quantum Monte Carlo computations of the team, have completed the half-a-century pursuit of the topological KT phase in quantum magnetic material, which eventually leads to the Nobel Physics Prize of 2016. The research findings are recently published in renowned academic journal *Nature Communications*.

KT phase of TMGO is detected

Quantum materials are becoming the cornerstone for the continuous prosperity of human society, including the next-generation AI computing chips that go beyond Moore's law, the high-speed Maglev train, and the topological unit for quantum computers, etc. However, these complicated systems require modern computational techniques and advanced analysis to reveal their microscopic mechanism. Thanks to the fast development of the supercomputing platforms all over the world, scientists and engineers are now making great use of these facilities to discover better materials that benefit our society. Nevertheless, computation cannot stand alone.

In the present investigation, experimental techniques for handling extreme conditions such as low temperature, high sensitivity and strong magnetic field, are required to verify the predictions and make discoveries. These equipments and technologies are acquired and organized by the team members coherently.

The research is inspired by the KT phase theory discovered by V Berezinskii, J Michael Kosterlitz and David J Thouless, of which the latter two are laureates of the Nobel Prize in Physics 2016 (together with F Duncan M Haldane) for their theoretical discoveries of topological phase, and phase transitions of matter. Topology is a new way of classifying and predicting the properties of materials, and now becoming the mainstream of quantum material research and industry, with broad potential applications in quantum computer,


lossless transmission of signals for information technology, etc. Back to 1970s, Kosterlitz and Thouless had predicted the existence of topological phase, hence named after them as the KT phase in quantum magnetic materials. Although such phenomena have been found in superfluids and superconductors, KT phase has yet been realized in bulk magnetic material, and is eventually discovered in the present work.

magnetic field and intrinsic magnetic moments of the material, which consequently allows the intricate topological KT fluctuations in the phase to be detected sensitively.

NMR spin-lattice relaxation rate measurements indeed revealed a KT phase sandwiched between a paramagnetic phase at temperature $T > T_u$ and an antiferromagnetic phase at temperature $T < T_l$.

This finding indicates a stable phase (KT phase) of TMGO, which serves as a concrete example of topological state of matter in crystalline material, might have potential applications in future information technologies. With its unique properties of topological excitations and strong magnetic fluctuations, many interesting research and potential applications with topological quantum materials can be pursued from here.

Dr. Meng said, "It will eventually bring benefits to the society, such that quantum computers, lossless transmission of signals for information technology, faster and more energy-saving high-speed trains, all these dreams could gradually come true from quantum material research."



NMR spectra and spin-lattice relaxation rates of TMGO in (a), (b) and (c) and its theoretical computation through large-scale QMC simulations in (d). Credit: The University of Hong Kong

To detect such interesting KT phase in a magnetic material is not easy, as usually the 3-dimensional coupling would render magnetic material to develop ordered phase but not topological phase at low temperature, and even if there exists a temperature window for the KT phase, highly sensitive measurement technique is required to be able to pick up the unique fluctuation pattern of the topological phase, and that is the reason why such phase has been enthusiastically perused, but its experimental discovery has defied many previous attempts. After some initial failures, the team member discovered that the NMR method under in-plane magnetic fields, do not disturb the low-energy electronic states as the in-plane moment in TMGO is mostly multipolar with little interference on

"Our approach, combining the state-of-art experimental techniques with unbiased quantum many-body computation schemes, enables us to directly compare experimental data to accurate numerical results with key theoretical predictions quantitatively, providing a bridge way to connect theoretical, numerical and experimental studies, the new paradigm set up by the joint team will certainly lead to more profound and impactful discoveries in quantum materials." He added.

The supercomputers used in computations and simulations

The powerful supercomputers Tianhe-1 and Tianhe-2 in China used in the computations are among the world's fastest supercomputers and ranked No.1 in 2010 and 2014 respectively in the TOP500 list (www.top500.org/). Their next-generation Tianhe-3 is expected to be in usage in 2021 and will be world first exaFLOPS scale supercomputer. The quantum Monte Carlo and tensor network simulations performed by the joint

team make use of the Tianhe supercomputers and requires the parallel simulations for thousands of hours on thousands of CPUs, it will take more than 20 years to finish if performed in common PC.

More information: Ze Hu et al. Evidence of the Berezinskii-Kosterlitz-Thouless phase in a frustrated magnet, *Nature Communications* (2020).
[DOI: 10.1038/s41467-020-19380-x](https://doi.org/10.1038/s41467-020-19380-x)

Provided by The University of Hong Kong

APA citation: Confirming simulated calculations with experiment results (2020, November 19) retrieved 27 January 2021 from <https://phys.org/news/2020-11-simulated-results.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.