

# Cattle grazing and soybean yields

January 6 2021

---



Little corn residue remains after concentrated grazing in the high stocking density treatment the day that cattle were moved from the fields in March of 2020. Credit: Morgan Grabau

By late fall, much of the Midwest is a pleasing landscape of dry, harvested corn fields. It makes for a bucolic rural scene on highway drives. But the corn litter that's left over doesn't seem useful, at least to untrained eyes.

But to those in the know, that corn residue is a valuable resource. Scattered leaves, husks, kernels, and cobs can serve as food to grazing cattle. When managed well, corn residue can increase farm income, provide affordable food for cattle, and efficiently use the land to feed people.

Morgan Grabau, a member of the American Society of Agronomy, studies the interactions of cattle grazing and crop productivity. She recently presented her research at the virtual ASA-CSSA-SSSA Annual Meeting.

"Corn residue is an under-used resource. Only 15% of the corn residue acres in the central U.S. are grazed," says Grabau.

One big concern farmers have about cattle grazing corn residue is [soil compaction](#). If cattle compact the [soil](#) too much, future crops might not grow well. Addressing the issue of soil compaction is the main focus of Grabau's work.

In the past, Grabau's research team has shown that compaction isn't too bad during fall and winter grazing. When the soil is dry and frozen, it resists stamping cattle hooves. "My research was focused on the effect of grazing in the spring when the soil is thawed and wet," she explains.



The normal stocking density treatment involved eight cattle grazing corn fields for 45 days. Credit: Morgan Grabau

Grabau studied two different grazing systems. In one system, researchers let a small number of cattle graze corn fields for 45 days starting in mid-February. The other system tripled the number of cattle but cut grazing time to just 15 days in March. This way, the total amount of grazing was equal. But the time spent on wet fields varied, which could affect how the soil responds to all that trampling.

The researchers studied corn fields in Nebraska, where around half of the corn fields are grazed after harvest. The team measured various soil

properties that contribute to compaction and the yield of the soybeans planted in the fields the following season after cattle were done grazing. The team repeated the experiment over two years.

"Much like previous fall grazing studies, minimal effects were seen on soil properties and yield due to spring grazing, regardless of the number of cattle and area grazed," says Grabau.

The soybean productivity of the fields following grazing did show some changes. The highly concentrated grazing for just 15 days actually increased yields slightly.

"This yield increase could be due to more residue removed, causing warmer soil temperatures for plants to grow," Grabau says.

The cattle did cause some soil compaction. But their effects were limited to the surface level of fields.



The high stocking density treatment had 24 cattle grazing corn fields for 15 days. This treatment increased future soybean yields slightly, perhaps because removing the corn residue warmed the soil. Credit: Morgan Grabau

"Compaction isn't permanent," Grabau says. "Soil can loosen up again as it dries and saturates over and over, and microbial activity in the soil also reduces compaction."

Fortunately, soybean seedlings had no problem establishing themselves in the soil after grazing even with some surface compaction present.

"Even when we created a [worst-case scenario](#), grazing in the spring when

the ground was wet, compaction was minimal and subsequent soybean yields were not negatively affected," Grabau says.

Although Grabau says that fall and winter grazing is probably still the best solution, farmers shouldn't be afraid of grazing cattle in the spring.

"The integration of crops and livestock is a beneficial production system," says Grabau. "Grazing [cattle](#) on corn residue can be a great way to make even more food for human consumption from corn fields, as both the [corn](#) grain and plant [residue](#) can be used as feed for livestock."

Provided by American Society of Agronomy

Citation: Cattle grazing and soybean yields (2021, January 6) retrieved 23 April 2024 from <https://phys.org/news/2021-01-cattle-grazing-soybean-yields.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.