

Red and green snow algae increase snowmelt in the Antarctic Peninsula

13 January 2021, by Kelsey Simpkins



Dr. Alia Khan and Chilean colleague Edgardo Sepulveda collect spectra albedo measurements in front of Collins Glacier on King George Island. Credit: Gonzalo Barrera

Red and green algae that grow on snow in the Antarctic Peninsula (AP) cause significant extra snowmelt on par with melt from dust on snow in the Rocky Mountains, according to a first-of-its-kind scientific research study led by Alia Khan, affiliate research scientist at the National Snow and Ice Data Center (NSIDC) and assistant professor at Western Washington University. Algal blooms are likely to increase in Antarctica as the planet continues to warm, which will further exacerbate seasonal snowmelt and contribute to the expansion of ice-free areas in the AP region. This could have serious impacts on regional climate, snow and ice melt, freshwater availability and ecosystems, yet is not accounted for in current global climate models. Results of the research were published on January 13, 2021, in the European Geosciences Union's *The Cryosphere*.

"Warming along the Antarctic Peninsula is causing drastic changes in snow and ice melt, as well as ecosystem responses," said Khan. "We are seeing these [algae](#) blooms spread across wide areas

along the coast. The blooms can be so intense and dark, like wearing a dark T-shirt on a sunny day, that they warm up the surface and cause more melting. The warming is likely expanding and strengthening the snow algae bloom season, which could continue to increase in this region of Antarctica as the climate continues to warm."

The researchers investigated the impacts of red and [green algae](#) on albedo, which is how much light the surface of the snow reflects back to space, and radiative forcing, which is how much energy the surface absorbs. Darker surfaces decrease albedo and increase radiative forcing, and positive radiative forcing causes the planet to warm. This is the first time that radiative forcing effects from algae have been estimated in the AP region. The scientists conducted this research by taking ground-based spectral albedo measurements of red and green snow algae at three sites in January 2018 and then modeled the radiative forcing using historical multi-year measurements of solar radiation at Palmer Station in Antarctica. Two of the ground sites were on King George Island, north of Palmer Station, and one was on Nelson Island. Because algal blooms are linked to animal waste, which produces the nutrients that the algae need to grow, the researchers chose sites that seals, penguins and other birds frequent.



Red and green snow algae in the Antarctic Peninsula region. Credit: Bob Gilmore

some unwanted consequences. We're seeing that these new darker and more widespread blooms are having a knock-on effect of accelerating melting, a feedback that makes the ice retreat a little faster."

The results of this research show that [snow](#) algae play a significant role in snowmelt in the AP regions where they occur, and should be accounted for in future estimates of ice-free expansion in the AP region. Future research will focus on a wider mapping the spread of [algae blooms](#), and on the climate events that create the most intense blooms.

More information: *The Cryosphere* (2021). DOI: [10.5194/tc-15-133-2021](https://doi.org/10.5194/tc-15-133-2021)

Provided by University of Colorado at Boulder

When compared with clean snow, the scientists found that green algae patches reduce snow albedo by 40 percent, and red algae patches reduce albedo by 20 percent. Green snow algae contain more chlorophyll than red snow algae and therefore absorb more solar radiation, reducing albedo by a greater amount for the same concentration of algae in the snow. As a result, radiative forcing averages are twice as high for green patches compared with red patches. During peak growing season, in austral summer, the increase in radiative forcing is about 26 watts per square meter, and about 13 watts per square meters for red algae. These amounts of [radiative forcing](#) are comparable to those caused by dust on snow in the mid-latitudes, such as in the Rocky Mountains, which advances snowmelt there. In the AP, [algal blooms](#) result in about 3,700 cubic meters, or over a million gallons of additional snow melt per year.

"Antarctica is already an incredibly beautiful place, and snow algae blooms add an extra artistic dimension," said Ted Scambos, a polar researcher at the Cooperative Institute for Research in the Environmental Sciences (CIRES) at the University of Colorado Boulder and co-author of the study. "But like so many natural systems, the change in climate is pushing things to new extremes, with

APA citation: Red and green snow algae increase snowmelt in the Antarctic Peninsula (2021, January 13) retrieved 25 January 2021 from <https://phys.org/news/2021-01-red-green-algae-snowmelt-antarctic.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.