


Study find physical weathering of rock breakdown more important than previously recognized

13 January 2021



Brad Carr, a University of Wyoming associate research scientist in geology and geophysics, uses a Geoprobe instrument to sample the subsurface in the foothills of the southern Sierra Nevada in California. Carr contributed to a study that was published in the Jan. 12 issue of *Geology*. The research shows that physical weathering is far more important than previously recognized in the breakdown of rock in mountain landscapes. Credit: Sarah Granke

Research led by the University of Wyoming shows that physical weathering is far more important than previously recognized in the breakdown of rock in mountain landscapes. Because it is difficult to measure, physical weathering has commonly been assumed to be negligible in previous studies.

Cliff Riebe, a professor in UW's Department of Geology and Geophysics, headed a research group that discovered that climate and erosion rates strongly regulate the relative importance of subsurface physical and chemical [weathering](#) of saprolite, the zone of weathered rock that retains the relative positions of mineral grains of the parent bedrock and lies between the layer of soil and

harder rock underneath. Saprolite is much like the weathered granite found on the flat areas surrounding the hard granite of Vedauwoo.

"Our work shows that physical strain can no longer be ignored in studies of subsurface weathering. It's not just a chemical process. It is physical as well," Riebe says. "What we found is that anisovolumetric weathering is much more common than previously thought, and that variations in this process can be explained by climate and erosion."

Riebe is lead author of a paper, titled "Anisovolumetric Weathering in Granitic Saprolite Controlled by Climate and Erosion Rates," which was published in the Jan. 12 issue of *Geology*. The journal publishes timely, innovative and provocative articles relevant to its international audience, representing research from all fields of the geosciences.

The study looked at three sites—with differing climates and elevations of granitic bedrock—of the Sierra Nevada, a mountain range in California.

In the lingo of geochemists, weathering has long been assumed to be "isovolumetric," meaning without a change in volume caused by physical strain.

"Our work shows that, to the contrary, weathering is commonly 'anisovolumetric,' meaning that strain caused by physical weathering is important," Riebe says.

Riebe credits some of the tools and instruments that were purchased from the Wyoming Center for Environmental Hydrology and Geophysics (WyCEHG) EPSCoR (Established Program to Stimulate Competitive Research) project that ended a few years ago as the reason his team could

measure both physical and chemical weathering at several sites in California.

"The reason why weathering was difficult to measure in the past is you have to be able to access the deep subsurface and sample it without disturbing it," Riebe explains. "You need a Geoprobe push coring system, which is basically a big track-mounted drill rig, to do this.

"It's expensive work, especially if you do not happen to own a Geoprobe and have to hire someone to do the work," he continues.

"Fortunately, we have access to this equipment and the expertise to operate it through Wyoming's Near Surface Geophysics facility, which is ably managed by Brad Carr, one of the study's co-authors."

The research was funded by grants from the National Science Foundation (NSF), NASA and the Natural Sciences and Engineering Research Council of Canada.

Riebe says there is a direct correlation between the research in this paper and the \$5.33 million NSF grant he received last September. The grant focuses on connections among rock, water and life at Earth's surface.

"This research is partly supported by that grant and also helped inspire it," Riebe says.

More information: Clifford S. Riebe et al, Anisovolumetric weathering in granitic saprolite controlled by climate and erosion rates *Geology* (2021) doi.org/10.1130/G48191.1

Provided by University of Wyoming

APA citation: Study find physical weathering of rock breakdown more important than previously recognized (2021, January 13) retrieved 17 January 2021 from <https://phys.org/news/2021-01-physical-weathering-breakdown-important-previously.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.