

# Potentially toxic plankton algae may play a crucial role in the future Arctic

February 3 2021

---



When the spring sets in in the Arctic, the metre-thick sea ice begins to melt. Melt ponds on the surface of the sea ice bring so much sunlight into the underlying seawater that the mixotrophic plankton algae start to grow dramatically. During an approx. 9-day period, the plankton can produce up to half of the total annual pelagic production in the high-Arctic fjord, Young Sound, in northeast Greenland. Several mixotrophic algae species are toxic. Credit: Lars Chresten Lund Hansen and Dorte H. Søgaard

As the sea ice shrinks in the Arctic, the plankton community that produces food for the entire marine food chain is changing. New research shows that a potentially toxic species of plankton algae that lives both via photosynthesis and absorbing food may become an important player in the Arctic Ocean as the future sea ice becomes thinner.

Microscopic [plankton algae](#), invisible to the naked eye, are the foundation of the marine food web, feeding all the ocean's living creatures from small crustaceans to large whales. Plankton algae need [light](#) and nutrients to produce food by photosynthesis.

A thick layer of sea ice—sometimes covered with snow—can reduce how much sunlight penetrates into the water and stop the algae getting enough light. However, as the sea ice is becoming thinner and less widespread in the Arctic, more light is penetrating into the sea. Does this mean more plankton algae and thus more food for more fish, whales and seabirds in the Arctic? The story is not so simple.

More light in the sea will only lead to a higher production of plankton algae if they also have enough nutrients—and this is often not the case. With the recent increase in freshwater melt from Arctic glaciers and the general freshening of the Arctic Ocean, more and more fresh and nutrient-depleted water is running out into the fjords and further out into the sea. The fresher water lies on top of the more salty ocean and stops nutrients from the deeper layers from mixing up towards the surface where there is light. And it is only here that plankton algae can be active.

## **Mixotrophic algae play on several strings**

However, a new study published today in the renowned journal *Scientific Reports* shows that so-called mixotrophic plankton algae may play a crucial role in the production of food in the Arctic Sea.

Mixotrophic algae are small, single-celled plankton algae that can perform photosynthesis but also obtain energy by eating other algae and bacteria. This allows them to live and grow even when there are insufficient light and nutrients for photosynthesis.



When the spring sets in in the Arctic, the metre-thick sea ice begins to melt. Melt ponds on the surface of the sea ice bring so much sunlight into the underlying seawater that the mixotrophic plankton algae start to grow dramatically. During an approx. 9-day period, the plankton can produce up to half of the total annual pelagic production in the high-Arctic fjord, Young Sound, in northeast Greenland. Several mixotrophic algae species are toxic. Credit: Lars Chresten Lund Hansen and Dorte H. Søgaard

In northeast Greenland, a team of researchers measured the production of plankton algae under the sea ice in the high-Arctic fjord Young Sound, located near Daneborg.

"We showed that the plankton algae under the sea ice actually produced up to half of the total annual plankton production in the fjord," says Dorte H. Søgaard from the Greenland Climate Research Centre, Greenland Institute of Natural Resources and the Arctic Research Centre, Aarhus University, who headed the study.

"Mixotrophic plankton algae have the advantage that they can sustain themselves by eating other algae and bacteria as a supplement to photosynthesis when there isn't enough light. This means that they are ready to perform photosynthesis even when very little light penetrates into the sea. In addition, many mixotrophic algae can live in relatively fresh water and at very low concentrations of nutrients—conditions that often prevail in the water layers under the sea ice in the spring when the ice melts," Dorte H. Søgaard explains.

## **Toxic algae kill fish**

For nine days, the researchers measured an algal bloom driven by mixotrophic algae occurring under the thick but melting sea ice in Young Sound during the Arctic spring in July, as the sun gained more power and more melt ponds spread across the sea ice, gradually letting through more light.

The algae belong to a group called haptophytes. Many of these algae are toxic, and in this study they bloomed in quantities similar to those previously observed in the Skagerrak near southern Norway. Here, the toxic plankton algae killed large amounts of salmon in Norwegian fish farms.

"We know that haptophytes often appear in areas with low salinity—as seen in the Baltic Sea, for example. It is therefore very probable that these mixotrophic-driven algae blooms will appear more frequently in a more freshwater-influenced future Arctic Ocean and that this shift in dominant algae to a mixotrophic algae species might have a large ecological and socio-economic impact," says Dorte H. Sjøgaard.

The researchers behind the project point out that it is the first time that a bloom of mixotrophic algae has been recorded under the sea ice in the Arctic.

**More information:** "An under-ice bloom of mixotrophic haptophytes in low nutrient and freshwater-influenced Arctic waters" *Scientific Reports* (2021). [doi.org/10.1038/s41598-021-82413-y](https://doi.org/10.1038/s41598-021-82413-y)

Provided by Aarhus University

Citation: Potentially toxic plankton algae may play a crucial role in the future Arctic (2021, February 3) retrieved 25 April 2024 from <https://phys.org/news/2021-02-potentially-toxic-plankton-algae-crucial.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.