

Nanomedicine activation profile determines efficacy depending on tumor c-Myc expression

26 February 2021




Fig. 1: Different drug release profile depending on the linker used for block-copolymers of nano-micelles FR-JQ1H/m with aliphatic aldehyde linker: Fast drug release in gradually elevated acidity SR-JQ1H/m with aromatic aldehyde linker: Slow drug release in gradually elevated acidity Drug release curve of FR-JQ1 and SR-JQ depending on the pH change of tumor tissues Credit: 2021 Innovation. Center of NanoMedicine

The Innovation Center of NanoMedicine reported in *ACS Nano* together with the group of Prof. Yu Matsumoto of Otorhinolaryngology and Head and Neck Surgery and the group of Prof. Horacio Cabral of the Department of Bioengineering in the University of Tokyo that the efficacy of polymeric nano-micelles with different drug activation profile depends on the expression level of c-Myc, one of the major proto-oncogenes, has been developed.


It is known that c-Myc is involved in cancer cell proliferation and angiogenesis and changes the cell cycle, suppresses normal cell differentiation, and promotes cancer metastasis. It is a typical proto-oncogene that regulates many genes related to growth factors and is known to be involved in developing of many cancers, such as

chromosomal translocation in Burkitt lymphoma. Therefore, drug discovery research is being conducted worldwide as an anticancer drug targeting this transcription factor that can directly attack cancer stem cells. However, since embryonic lethality occurs in c-Myc knockout mice, c-Myc is considered as an essential gene for living cells, and selective delivery to cancer tissues is an important key to developing its inhibitors. Besides, c-Myc is also known as a factor necessary for the initial induction of iPS cells. In the future this inhibition can be expected to be applied as a technology that can also be used to suppress iPS cell-derived carcinogenesis.

In this study, JQ1H, which is a structural analogue of JQ1, a typical indirect c-Myc inhibitor, was encapsulated inside functional nano-micelles, and their efficacy was evaluated. JQ1 binds to a bromodomain protein called BRD4, which is involved in the activation of RNA polymerase II regulating the expression of c-Myc, to inhibit this stream strongly. As a result, the activity of RNA polymerase is weakened and c-Myc expression is down-regulated. Although JQ1 was expected as a promising epigenome drug due to its strong gene expression inhibition, it has an extremely short half-life in vivo due to its fast kidney excretion and rapid clearance after administration. Additionally, JQ1 is almost insoluble in water. These properties of JQ1 became big issues to develop it as an effective drug. The polymeric nano-micelles developed so far at the Innovation Center of NanoMedicine (iCONM), for anticancer therapy, demonstrated (1) stabilization of encapsulated drugs, (2) suppression of kidney excretion, (3) EPR (selective drug delivery to cancer tissues) mediated tumor accumulation, and (4) drug release based on tumor acidosis. This time, we confirmed good antitumor activity in mice transplanted with tongue cancer, melanoma and pancreatic cancer using


JQ1-equipped nano-micelles.

copolymer micelles in nanomedicine applications" Chem. Rev. 2018, 118 6844-6892. (DOI: 10.1021/acs.chemrev.8b00199). Credit: 2021 Innovation Center of NanoMedicine



upper: tongue cancer, lower: pancreatic cancer. Credit: 2021 Innovation Center of NanoMedicine

Nano-micelles containing JQ1H leak into the tumor tissue from blood vessels after systemic administration due to the so-called EPR effect. Tumor tissues are rich in [lactic acid](#) due to its enhanced glycolysis and is more acidic than normal tissues. In this work, two types of nano-micelle were prepared; one in which hydrophobic JQ1H were linked to a amphiphilic block polymer composed of hydrophilic polyethylene glycol block and hydrophobic poly-amino acid block using 3-aminopropionaldehyde (aliphatic aldehyde) linker and the other micelle in which JQ1H was linked with polymer via p-aminomethylbenzaldehyde (aromatic aldehyde) linker. An amphiphilic block polymer was synthesized and used as a base material for nano-micelles. When it was self-assembled in water to a micellar structure and administered to cancer-bearing mice, antitumor activity was achieved.



When the linker is an aliphatic aldehyde or when it is an aromatic aldehyde, the release pattern of the drug differs greatly depends on the acidity. The former releases the drug rapidly, and the latter releases the [drug](#) slowly. Therefore, the former nano-medicine was named FR-JQ1H/m and the latter was named SR-JQ1H/m. The antitumor activity of these nano-micelles differ greatly depending on the expression level of c-Myc. While, FR-JQ1H/m is more effective for tumors with high c-Myc expression, SR-JQ1H/m is more effective for tumors with low c-Myc expression.

Polymeric micelles were one of the first polymer self-assemblies reported as a nano-DDS, and are composed of distinct two domains, a drug-loading core and a hydrophilic shell. Amphiphilic block copolymers, containing a hydrophilic block and a hydrophobic block, are firstly revealed to construct those distinct domains in a micelle structure through spontaneous self-assemble as a result of hydrophobic interactions in aqueous. H. Cabral, K. Miyata, K. Osada, K. Kataoka, "Block

In the future, we believe that the selection of nano-micelles according to the expression level of biomarkers will be an important step toward the realization of personalized medicine and in-body hospitals.

More information: Hitoshi Shibasaki et al, Efficacy of pH-Sensitive Nanomedicines in Tumors with Different c-MYC Expression Depends on the Intratumoral Activation Profile, *ACS Nano* (2021).

[DOI: 10.1021/acsnano.1c00364](https://doi.org/10.1021/acsnano.1c00364)

Provided by Innovation Center of NanoMedicine

APA citation: Nanomedicine activation profile determines efficacy depending on tumor c-Myc expression (2021, February 26) retrieved 15 May 2021 from <https://phys.org/news/2021-02-nanomedicine-profile-efficacy-tumor-c-myc.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.