


Information transition mechanisms of spatiotemporal metasurfaces

11 March 2021



a, The conceptual illustration of manipulating the transmission of electromagnetic waves via spatiotemporal metasurface. b, The input phase states of the meta-atom with $N=4$. c, Cayley diagram of the product group $Z_N \times Z_q$ ($Z_4 \times Z_3$) that generates $N \times q$ (4×3) output phase states. d, The generated output spectral states of each meta-atom e, The schematic of independent control of spectral responses of the meta-atom at two harmonics, in which each line connects the spectral responses of the meta-atom generated by the same temporal sequence. Credit: Haotian Wu, Xin Xin Gao, Lei Zhang, Guo Dong Bai, Qiang Cheng, Lianlin Li, and Tie Jun Cui

information perspective, in which two information transition mechanisms on group extension and independent control of multiple harmonics are revealed and characterized. The information transition efficiencies of these mechanisms are analyzed as well, which could be used to predict the channel capacity of the spatiotemporal metasurfaces for wireless communications. The presented framework and gained results would be helpful to lay the groundwork for information-based spatiotemporal metasurfaces.

Spatiotemporal metasurfaces, driven by ultrafast dynamic modulations, have opened up new possibilities for manipulating the harmonic modes of electromagnetic waves and generations of exotic physical phenomena, such as dispersion cancelation, Lorentz reciprocity breaking, and Doppler illusions. In recent years, rapid development of information technologies have stimulated many information processing applications for metasurfaces, including computational imaging, wireless communications, and performing mathematical operations.

With increasing research focused on the topic of information processing with metasurfaces, a general theory is urgently required to characterize the information processing abilities of the spatiotemporal metasurfaces. In a new paper published in *Light Science & Applications*, Prof. Tie Jun Cui's group at Southeast University (SEU) has reported a breakthrough on this topic. In this work, the information transition mechanisms of spatiotemporal metasurfaces are proposed and analyzed, in which the group extension and independent control of multiple harmonics are revealed and characterized as two major information transition mechanisms of the spatiotemporal metasurfaces.

Specifically, the group extension mechanism could be adopted to extend the output phase states of each meta-atom by a factor of q , where q is a

Spatiotemporal metasurfaces are analyzed from an

function over the modulation periodicity, input phase information-oriented applications including cognitive states, and harmonic index. Accordingly, the output harmonic wavefront engineering, intelligent spectral response states of the spatiotemporal computational imaging, and the 6th generation (6G) [metasurface](#) are extended greatly, such that more accurate manipulations of the electromagnetic wireless communications," the scientists conclude. information could be obtained without increasing the design complexity of the metasurfaces.

Additionally, the researchers demonstrated that the independent control of the spectral responses of the spatiotemporal metasurface could be realized as well. The independent control of multiple harmonics could open up new possibilities for the metasurface-based multitasking, by which electromagnetic information could be independently processed with frequency-gapped channels. A proof-of-concept experiment is performed in the microwave regime to verify the mechanisms of the group extension and independent control of multiple harmonics with the spatiotemporal metasurface.

By incorporating the proposed model with the Shannon's entropy theory, the authors further discovered the information transition efficiencies of the spatiotemporal metasurfaces with respect to the above two mechanisms. The obtained results could be applied to predict the channel capacity of the spatiotemporal metasurface, which would be helpful to guide the analysis and design of spatiotemporal metasurfaces for wireless communications. Moreover, they demonstrated that the output spectral responses of the spatiotemporal metasurface can help prove Fermat's little theorem, which in turn might provide more clues to understand the non-vanished spectra responses of the spatiotemporal metasurfaces.

"The proposed theory establishes a quantitative framework to characterize the information transition capabilities of the spatiotemporal metasurfaces, which provides deeper physical insights in understanding spatiotemporal metasurfaces from the information perspective, and offers new approaches to facilitate analysis and design. The presented framework and obtained results, with wide-ranging spectral applicability, would be helpful to lay the groundwork for future researches into the regime of information-based spatiotemporal metasurfaces, and expected to enable new

More information: Haotian Wu et al, Harmonic information transitions of spatiotemporal metasurfaces, *Light: Science & Applications* (2020). [DOI: 10.1038/s41377-020-00441-1](https://doi.org/10.1038/s41377-020-00441-1)

Provided by Chinese Academy of Sciences

APA citation: Information transition mechanisms of spatiotemporal metasurfaces (2021, March 11)
retrieved 17 June 2021 from <https://phys.org/news/2021-03-transition-mechanisms-spatiotemporal-metasurfaces.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.