


Algorithms inspired by social networks reveal lifecycle of substorms, a key element of space weather

March 23 2021



Map representing a snapshot of the community structure at onset. Credit: Background map credit: SuperMAG

Space weather often manifests as substorms, where a beautiful auroral display such as the Northern Lights is accompanied by an electrical

current in space which has effects at earth that can interfere with and damage power distribution and electrical systems. Now, the lifecycle of these auroral substorms has been revealed using social media-inspired mathematical tools to analyse space weather observations across the Earth's surface.


Analysis by researchers led by the University of Warwick has revealed that these substorms manifest as global-scale [electrical current](#) systems associated with the spectacular aurora, reaching across over a third of the globe at [high latitudes](#).

New research which involves the University of Warwick, John Hopkins University—Applied Physics Laboratory, University of Bergen and Cranfield University, and published today in the journal *Nature Communications* processes data on disturbances in the Earth's [magnetic field](#) from over a hundred magnetometers in the Northern hemisphere using a new technique that enables them to find 'like-minded friends'.

Magnetometers register changes in the Earth's magnetic field. When charged particles from our Sun bombard the Earth's magnetic field, it stores up energy like a battery. Eventually, this energy is released leading to large-scale electrical currents in the ionosphere which generate disturbances of magnetic fields on the ground. At extremes, this can disrupt power lines, electronic and [communications systems](#) and technologies such as GPS.

Using [historical data](#) from the SuperMAG collaboration of magnetometers, the researchers applied algorithms from network science to find correlations between magnetometer signals during 41 known substorms that occurred between 1997-2001. These use the same principles that allow a social networking site to recommend new friends, or to push relevant advertisements to you as you browse the internet.

Magnetometers detecting coherent signals were linked into communities, regardless of where they were located on the globe. As time progressed, they saw each substorm develop from many smaller communities into a single large correlated system or community at its peak. This led the authors to conclude that substorms are one coherent current system which extends over most of the nightside high latitude globe, rather than a number of individual small and disjointed current systems.



Map representing a snapshot of the community structure at the time of maximum auroral expansion. Credit: Background map credit: SuperMAG

Dr. Lauren Orr, who led the research as part of her Ph.D. at the University of Warwick Department of Physics and is now based at

Lancaster University, said: "We used a well-established method within [network science](#) called community detection and applied it to a [space weather](#) problem. The idea is that if you have lots of little subgroups within a big group, it can pick out the subgroups.

"We applied this to space weather to pick out groups within magnetometer stations on the Earth. From that, we were trying to find out whether there was one large current system or lots of separate individual current systems.

"This is a good way of letting the data tell us what's going on, instead of trying to fit observations to what we think is occurring."

Some recent work has suggested that auroral substorms are composed of a number of smaller electrical current systems and remain so throughout their lifecycle. This new research demonstrates that while the substorm begins as lots of smaller disturbances, it quite rapidly becomes a large system over the course of around ten minutes. The lack of correlation in its early stages may also suggest that there is no single mechanism at play in how these substorms evolve.

The results have implications for models designed to predict space weather. Space [weather](#) was included in the UK National Risk Register in 2012 and updated in 2017 with a recommendation for more investment in forecasting.

Co-author Professor Sandra Chapman adds: "Our research introduces a whole new methodology for looking at this data. We've gone from a data poor to a data rich era in space plasma physics and [space weather](#), so we need new tools. It's a first to show that you can take one of these tools to our field and get a really important result out of it. We've had to learn a lot to be able to do that, but in doing so it opens up a new window into the data."

More information: "Network Community Structure of Substorms using SuperMAG Magnetometers" *Nature Communications*, [DOI: 10.1038/s41467-021-22112-4](https://doi.org/10.1038/s41467-021-22112-4)

Provided by University of Warwick

Citation: Algorithms inspired by social networks reveal lifecycle of substorms, a key element of space weather (2021, March 23) retrieved 20 September 2024 from <https://phys.org/news/2021-03-algorithms-social-networks-reveal-lifecycle.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.