

# Study shows promise of forecasting meteotsunamis

April 1 2021

---



The Lake Michigan meteotsunami unleashes a waterfall over the breakwater during the event near the Ludington, Michigan, Breakwater Lighthouse on April 13, 2018. Credit: Debbie Maglothin

On the afternoon of April 13, 2018, a large wave of water surged across

Lake Michigan and flooded the shores of the picturesque beach town of Ludington, Michigan, damaging homes and boat docks, and flooding intake pipes. Thanks to a local citizen's photos and other data, NOAA scientists reconstructed the event in models and determined this was the first ever documented meteotsunami in the Great Lakes caused by an atmospheric inertia-gravity wave.

An atmospheric inertia-gravity wave is a wave of air that can run from 6 to 60 miles long that is created when a mass of stable air is displaced by an air mass with significantly different pressure. This sets in motion a wave of air with rising and falling pressure that can influence the [water](#) below, as it synchronizes with water movement on the [lake](#)'s surface like two singers harmonizing.

"That meteotsunami was hands down off the chart awesome," said Debbie Maglothin of Ludington who took photos of the event. "The water in between the breakwaters didn't go down like the water on the outside of them, so it created waterfalls that cascaded over the breakwaters. Had this event occurred during summer it could have washed people right off the breakwaters."

Meteotsunamis generated from this type atmospheric condition are common around the globe, but in the Great Lakes, the few well documented meteotsunamis have been driven by sudden [severe thunderstorms](#) where both winds and air pressure changes have played significant roles.

## **Combining water and weather models**

While there are currently no forecast models that effectively predict meteotsunamis in the U.S., new NOAA research based on the Ludington wave demonstrates that existing NOAA numerical weather prediction models and hydrodynamic forecast models may enable scientists to

predict these meteotsunami-driving atmospheric waves minutes to hours in advance. The research is published in a special edition of the journal *Natural Hazards* about meteotsunamis.

"The good news with this type of meteotsunami is that it is easier to predict than ones triggered by thunderstorms," said Eric Anderson, an oceanographer at NOAA's Great Lakes Environmental Research Laboratory and lead author of the study. "Our short-range [weather models](#) can pick up these atmospheric pressure waves, whereas predicting thunderstorms is more difficult."

Meteotsunamis are a lesser known category of tsunami. Unlike the more well known tsunami—such as the catastrophic 2004 Boxing Day tsunami in Indonesia, which was caused by an earthquake on the seafloor, meteotsunamis are caused by weather, in particular some combination of changing air pressure, [strong winds](#) and thunderstorm activity.

"Because the lakes are relatively small, meteotsunamis typically need more than a jump in air pressure to drive them," said Anderson. "That's where the thunderstorms and wind come in to give them a push."

## **Great Lakes have history of meteotsunamis**

Meteotsunamis occur around the world, and are known to occur in the United States primarily on the Great Lakes and along the East and Gulf of Mexico coasts. Meteotsunami waves in the Great Lakes can be particularly insidious because they can bounce off the shoreline and come back again when the skies are clear. They are relatively rare and typically small, the largest producing three to six foot waves, which only occur about once every 10 years.

Predicting these waves in advance would give communities potentially life-saving warnings and would allow residents and businesses to take

measures to better protect property. The Ludington meteotsunami resulted in some property damage but no serious injuries. Had the meteotsunami struck in the summer when swimmers, anglers and vacationers flock to the lakeshore beaches, parks and waters, it might have been a different story, as was the case with a meteotsunami that took the lives of eight people in Chicago in June 1954.

"It's a gap in our forecasting," said Anderson. "With this study and other research we are getting closer to being able to predict them in advance."

**More information:** Eric J. Anderson et al, A high-amplitude atmospheric inertia–gravity wave-induced meteotsunami in Lake Michigan, *Natural Hazards* (2020). [DOI: 10.1007/s11069-020-04195-2](https://doi.org/10.1007/s11069-020-04195-2)

Provided by NOAA Headquarters

Citation: Study shows promise of forecasting meteotsunamis (2021, April 1) retrieved 26 April 2024 from <https://phys.org/news/2021-04-meteotsunamis.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.