

# **Qubits composed of holes could be the trick to build faster, larger quantum computers**

April 2 2021

---



Credit: Pixabay/CC0 Public Domain

A new study indicates holes the solution to operational speed/coherence trade-off, potential scaling up of qubits to a mini-quantum computer.

Quantum computers are predicted to be much more powerful and functional than today's 'classical' computers.

One way to make a quantum bit is to use the 'spin' of an electron, which can point either up or down. To make quantum computers as fast and power-efficient as possible we would like to operate them using only electric fields, which are applied using ordinary electrodes.

Although spin does not ordinarily 'talk' to electric fields, in some materials spins can interact with electric fields indirectly, and these are some of the hottest materials currently studied in quantum computing.

The interaction that enables spins to talk to electric fields is called the [spin-orbit interaction](#), and is traced all the way back to Einstein's theory of relativity.

The fear of [quantum-computing](#) researchers has been that when this interaction is strong, any gain in operation speed would be offset by a loss in coherence (essentially, how long we can preserve quantum information).

"If electrons start to talk to the electric fields we apply in the lab, this means they are also exposed to unwanted, fluctuating electric fields that exist in any material (generically called 'noise') and those electrons' fragile quantum information would be destroyed," says A/Prof Dimi Culcer (UNSW/FLEET), who led the theoretical roadmap study.

"But our study has shown this fear is not justified."

"Our [theoretical studies](#) show that a solution is reached by using holes, which can be thought of as the absence of an electron, behaving like positively-charged electrons."

In this way, a quantum bit can be made robust against charge fluctuations stemming from the solid background.

Moreover, the '[sweet spot](#)' at which the [qubit](#) is least sensitive to such noise is also the point at which it can be operated the fastest.

"Our study predicts such a point exists in every [quantum bit](#) made of holes and provides a set of guidelines for experimentalists to reach these points in their labs," says Dimi.

Reaching these points will facilitate experimental efforts to preserve [quantum information](#) for as long as possible. This will also provide strategies for 'scaling up' quantum bits—ie, building an 'array' of bits that would work as a mini-quantum [computer](#).

"This theoretical prediction is of key importance for scaling up quantum processors and first experiments have already been carried out," says Prof Sven Rogge of the Centre for Quantum Computing and Communication Technology (CQC2T)."

"Our recent experiments on hole qubits using acceptors in silicon already demonstrated longer coherence times than we expected," says A/Prof Joe Salfi of the University of British Columbia. "It is encouraging to see that these observations rest on a firm theoretical footing. The prospects for hole qubits are bright indeed."

The paper, "Optimal operation points for ultrafast, highly coherent Ge hole [spin](#)-orbit qubits," was published in Nature partner journal *npj Quantum Information* in April 2021.

**More information:** Zhanning Wang et al. Optimal operation points for ultrafast, highly coherent Ge hole spin-orbit qubits, *npj Quantum Information* (2021). [DOI: 10.1038/s41534-021-00386-2](https://doi.org/10.1038/s41534-021-00386-2)

Provided by Australian Research Council

Citation: Qubits composed of holes could be the trick to build faster, larger quantum computers (2021, April 2) retrieved 21 September 2024 from <https://phys.org/news/2021-04-qubits-holes-faster-larger-quantum.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.