


Green chemistry and biofuel: The mechanism of a key photoenzyme decrypted

8 April 2021



Artist impression of the enzymatic catalysis proposed in mechanism of fatty acid photodecarboxylase. Credit: Damien Sorigué

The functioning of the enzyme FAP, useful for producing biofuels and for green chemistry, has been decrypted. This result mobilized an international team of scientists, including many French researchers from the CEA, CNRS, Inserm, École Polytechnique, the universities of Grenoble Alpes, Paris-Saclay and Aix Marseille, as well as the European Synchrotron (ESRF) and synchrotron SOLEIL. The study is published in *Science* on April 09, 2021.


The researchers decrypted the operating mechanisms of FAP (Fatty Acid Photodecarboxylase), which is naturally present in microscopic algae such as *Chlorella*. The enzyme had been identified in 2017 as able to use [light energy](#) to form hydrocarbons from [fatty acids](#) produced by these microalgae. To achieve this new result, research teams used a complete experimental and theoretical toolkit.

Understanding how FAP works is essential because this photoenzyme opens up a new opportunity for sustainable biofuel production from

fatty acids naturally produced by living organisms. FAP is also very promising for producing high added-value compounds for fine chemistry, cosmetics and pharmaceuticals.

In addition, due to their light-induced reaction, photoenzymes give access to ultrarapid phenomena that occur during enzymatic reactions. FAP therefore offers a unique opportunity to understand in detail a chemical reaction taking place in living organisms.

More specifically, in this work, researchers show that when FAP is illuminated and absorbs a photon, an electron is stripped in 300 picoseconds from the fatty acid produced by the algae. This fatty [acid](#) is then dissociated into a hydrocarbon precursor and carbon dioxide (CO₂). Most of the CO₂ generated is then turned in 100 nanoseconds into bicarbonate (HCO₃⁻) within the enzyme. This activity uses light but does not prevent photosynthesis: the flavin molecule within the FAP, which absorbs the photon, is bent. This conformation shifts the molecule's absorption spectrum towards the red, so that it uses photons not used for the microalgae's photosynthetic activity.



The researchers decrypted the operating mechanisms of FAP (Fatty Acid Photodecarboxylase), which is naturally present in microscopic algae such as *Chlorella*. Credit: Damien Sorigue

It is the combined interpretation of the results of various experimental and theoretical approaches by the [international consortium](#) that yields the detailed, atomic-scale picture of FAP at work. This multidisciplinary study combined bioengineering work, optical and [vibrational spectroscopy](#), static and kinetic crystallography performed with synchrotrons or an X-ray free electron laser, as well as quantum chemistry calculations.

More information: "Mechanism and dynamics of fatty acid photodecarboxylase" *Science* (2021). science.sciencemag.org/cgi/doi/10.1126/science.abd5687

Provided by European Synchrotron Radiation Facility

APA citation: Green chemistry and biofuel: The mechanism of a key photoenzyme decrypted (2021, April 8) retrieved 15 May 2021 from <https://phys.org/news/2021-04-green-chemistry-biofuel-mechanism-key.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.