

# Conservationists may be unintentionally spreading pathogens between threatened animal populations

April 12 2021

---



An at-risk mussel species. Credit: David Aldridge

Moving endangered species to new locations is often used as part of species conservation strategies, and can help to restore degraded

ecosystems. But scientists say there is a high risk that these relocations are accidentally spreading diseases and parasites.

The new report published today in the journal *Conservation Letters* focuses on [freshwater mussels](#), which the researchers have studied extensively, but is applicable to all [species](#) moved around for conservation purposes.

Mussels play an important role in cleaning the water of many of the world's rivers and lakes, but are one of the most threatened [animal groups](#) on Earth. There is growing interest in moving mussels to new locations to boost threatened populations, or so they can be used as 'biological filters' to improve water quality.

A gonad-eating [parasitic worm](#), *Rhipidocotyle campanula*, which can leave mussels completely sterile, was identified as a huge risk for captive breeding programmes where mussels from many isolated populations are brought together.

"We need to be much more cautious about moving animals to new places for conservation purposes, because the costs may outweigh the benefits," said Dr. David Aldridge in the Department of Zoology at the University of Cambridge, senior author of the report.

He added: "We've seen that mixing different populations of mussels can allow widespread transmission of gonad-eating worms—it only takes one infected mussel to spread this parasite, which in extreme cases can lead to collapse of an entire population."

Pathogens can easily be transferred between locations when mussels are moved. In extreme cases, the pathogens may cause a population of [mussels](#) to completely collapse. In other cases infections may not cause a problem unless they are present when other factors, such as lack of food

or high temperatures, put a population under stress leading to a sudden outbreak.

The report recommends that species are only relocated when absolutely necessary and quarantine periods, tailored to stop transmission of the most likely pathogens being carried, are used.

It identifies four key factors that determine the risk of spreading pathogens when relocating animals: proportion of infected animals in both source and recipient populations; density of the resulting [population](#); host immunity; and the life-cycle of the pathogen. Pathogens that must infect multiple species to complete their life-cycle, like parasitic mites, will only persist if all of the species are present in a given location.

"Moving animals to a new location is often used to protect or supplement endangered populations. But we must consider the risk this will spread [pathogens](#) that we don't understand very well at all, which could put these populations in even greater danger," said Josh Brian, a Ph.D. student in the Department of Zoology at the University of Cambridge and first author of the report.

Different populations of the same species may respond differently to infection with the same pathogen because of adaptations in their immune system. For example, a pack of endangered wolves moved to Yellowstone National Park died because the wolves had no immunity to parasites carried by the local canines.

The researchers say that stocking rivers with fish for anglers, and sourcing exotic plants for home gardens could also move around parasites or diseases.

"Being aware of the risks of spreading diseases between populations is a vital first step towards making sure we avoid unintentional harm in

future conservation work," said Isobel Ollard, a Ph.D. student in the Department of Zoology at the University of Cambridge, who was also involved in the study.

**More information:** *Conservation Letters* (2021). [DOI: 10.1111/conl.12799](https://doi.org/10.1111/conl.12799)

Provided by University of Cambridge

Citation: Conservationists may be unintentionally spreading pathogens between threatened animal populations (2021, April 12) retrieved 22 September 2024 from <https://phys.org/news/2021-04-conservationists-unintentionally-pathogens-threatened-animal.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--