


Researcher questions whether powered flight appeared on non-avian dinosaurs

26 April 2021



Credit: F.J. Serrano. University of Malaga

Powered flight in animals—that uses flapping wings to generate thrust—is a very energetically demanding mode of locomotion that requires many anatomical and physiological adaptations. In fact, the capability to develop it has only appeared four times in the evolutionary history of animals: On insects, pterosaurs, birds and bats.

A [research paper](#) published in 2020 in the scientific journal *Current Biology* concluded that, apart from birds, the only living descendants of dinosaurs, powered flight would have originated independently in other three groups of dinosaurs. This is a conclusion that makes a great impact, as it increases the number of vertebrates that would have developed this costly mode of locomotion, which, among dinosaurs, would no longer be an exclusive capability of birds.

The scientist of the Department of Ecology and Geology of the University of Malaga Francisco Serrano Alarcón has recently published an article in the same journal, questioning the idea that powered flight appeared multiple times among dinosaurs.

The researcher of the UMA, member of the Dinosaur Institute (NHMLAC) of Los Angeles, refutes such conclusion in the absence of scientific evidence. As he remarks, the parameters used by the authors to determine flight capability do not allow differentiation between powered flight and passive flight, the latter being frequent in many more animal groups.



Credit: Stephanie Abramowicz (Natural History Museum of Los Angeles County).

This new study, which he conducted along with the paleontologist Luis M. Chiappe, Vice-President for Research and Collections of the NHMLAC, compares the parameters measured on present animals with powered flight capability, such as [birds](#) and bats, and gliding [animals](#), for example, flying squirrels or flying reptiles, among others. Moreover, they added new data on the capability to generate energy from muscles in addition to the data considered in the original study.

"Birds are a group of dinosaurs of which we have discovered 150-million-year-old fossils with fully developed wings. Among their closest non-avian relatives, we have also found fossils with

sufficiently developed wings that could provide them with some aerodynamic benefit, whether to glide between trees or get thrust to climb and jump over obstacles. But this does not mean that they could take off by flapping their wings or maintain a powered flight," explains Francisco Serrano.

In short, both authors conclude that although they cannot discount the possibility that powered [flight](#) appeared in other non-avian [dinosaurs](#), current evidence does not support the hypothesis suggested in the original paper by Pei et al (2020).

More information: Francisco J. Serrano et al, Independent origins of powered flight in paravian dinosaurs?, *Current Biology* (2021). [DOI: 10.1016/j.cub.2021.03.058](#)

Provided by University of Malaga

APA citation: Researcher questions whether powered flight appeared on non-avian dinosaurs (2021, April 26) retrieved 22 June 2021 from <https://phys.org/news/2021-04-powered-flight-non-avian-dinosaurs.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.