


Previously unrecognized tsunami hazard identified in coastal cities

May 3 2021, by Lois Yoksoulian



Evolution of earthquake-induced tsunami in a bay traversed by a strike-slip fault. Colors indicate sea-surface height relative to undisturbed water level; wave

height may exceed 2 m in some places. Top: Instantaneous tsunami generation phase coincident with earthquake propagation; Middle: postseismic tsunami generation phase, in which gravity-driven tsunami waves dominate; Bottom: backward propagation of large wave fronts reflected from bay tip. Credit: Ares J. Rosakis

A new study found overlooked tsunami hazards related to undersea, near-shore strike-slip faults, especially for coastal cities adjacent to faults that traverse inland bays. Several areas around the world may fall into this category, including the San Francisco Bay area, Izmit Bay in Turkey and the Gulf of Al-Aqaba in Egypt.

The study led by University of Illinois Urbana-Champaign civil and environmental engineering professor Ahmed Elbanna and professor Ares Rosakis of the California Institute of Technology used the Blue Waters supercomputer at the National Center for Supercomputing Applications to model tsunami hazards related to strike-slip faults around the globe. The results are published in the *Proceedings of the National Academy of Sciences*.

"Whenever we saw large tsunamis triggered by earthquakes along strike-slip faults, people assumed that perhaps the [earthquake](#) had caused an undersea landslide, displacing water that way," Rosakis said.

The researchers said that a strike-slip [fault](#) exists when two blocks of rock on the fault line slide horizontally past one another. The San Andreas Fault is an example of a strike-slip fault.

In September 2018, a moderate 7.5 magnitude earthquake and unexpectedly powerful tsunami swept through Palu, a city situated on the inland side of Palu Bay on the Indonesian island of Sulawesi. The quake

occurred along a northwest-southeast trending strike-slip fault that runs through the city and plunges below the bay along Palu's northwest shore.

"It looked like a bulldozer had come in and leveled the town," said co-author Costas Synolakis, the president of Athens College and a professor of civil engineering at the University of Southern California, who surveyed the area following the devastating event. "This is why it is so important that we try to understand what really happened."

Studies exploring connections between strike-slip faulting and tsunamis exist. However, they focus on specific fault systems or geographic locations, obscuring the complex details of the fault geometry and bathymetry, the study reports.

"What is unique about our study is that instead of considering a location-specific event, we focused on the fundamentals of a strike-slip fault system interacting within the boundaries of a narrow bay," Elbanna said. "We opted to simulate a very basic planar fault passing through a very simplified smooth-bottomed bay, similar to a bathtub. Having this simplified baseline model allows us to generalize to any place on the planet that may be at risk."

Intersonic earthquakes are fault ruptures that happen so quickly that their movement outpaces the seismic shear waves they generate—like a sonic boom, but with the shock wave moving through the earth's crust. The simulations found that intersonic earthquakes can provide enough energy and large enough horizontal displacements to trigger large tsunami waves.

When such earthquakes occur within a narrow bay, the researchers reported three [distinct phases](#) that can lead to a tsunami: the initial fault movement and shockwave causing almost instantaneous shaking of the coastal land; the displacement of water while the earthquake is

occurring; and gravity-driven motion of the tsunami wave after the ground motion has subsided that carries the wave to shore.

"Each of these phases will have a different effect depending on the unique geography of the surrounding land and bathymetry of the bay," Elbanna said. "And, unlike the earthquakes and subsequent water displacement that occur many miles offshore, an earthquake and tsunami that occurs within the narrow confines of a bay will allow for very little warning time for the coast."

Elbanna compares the effect of horizontal strike-slip fault displacements to holding a water cup in your hand and shaking it horizontally.

"The sloshing motion is a result of the horizontal shaking. When an earthquake occurs along a strike-slip fault in a narrow bay, the horizontal ground motion pushes and pulls the boundaries of the bay leading to displacement of water in the vertical direction and initiation of the tsunami," he said.

"The physics-based model used in this study provides critical insight about the hazard associated with strike-slip faulting, particularly, the need to account for such risk to mitigate future damage to other bays traversed by strike-slip faults," said Illinois graduate student Mohamed Abdelmeguid, who conducted the simulations along with former graduate student Xiao Ma, currently a senior research scientist at Exxon Mobil.

The at-risk regions identified by the team—Northern California, Turkey and Egypt—have experienced intersonic earthquakes in the past, and the researchers recommend revisiting the tsunami hazard rating of underwater strike-slip faults, particularly those traversing narrow bays.

"It may not look like the tsunami scene from Dwayne Johnson's "San

Andreas' movie, but the [tsunami](#) risk for Northern California and several places worldwide needs to be seriously revisited," Elbanna said.

More information: Ahmed Elbanna et al., "Anatomy of strike-slip fault tsunami genesis," *PNAS* (2021).

www.pnas.org/cgi/doi/10.1073/pnas.2025632118

Provided by University of Illinois at Urbana-Champaign

Citation: Previously unrecognized tsunami hazard identified in coastal cities (2021, May 3)
retrieved 20 September 2024 from

<https://phys.org/news/2021-05-previously-unrecognized-tsunami-hazard-coastal.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.