

'Smart collar' could prevent tapeworms in dogs

July 8 2021



Development and field tests of smart deworming collar (A: 3D stacked graph of smart collar; B: Embedded modules for smart collar; C: Overall shape of Smart collar; D: Recovery of collars in July 2019 in Seni district after they had been attached for a year). Credit: Yang S-J et al., 2021, PLOS Neglected Tropical Diseases, CC-BY 4.0 (creativecommons.org/licenses/by/4.0/)

Dogs infected with echinococcosis play a major role in spreading tapeworms across human populations around the world. Now, researchers have developed a "smart collar" which gradually delivers a steady dose of a deworming drug to dogs. The collar successfully reduces the animals' risk of echinococcosis, the team reports in *PLOS Neglected Tropical Diseases*

Dogs can be infected with either *Echinococcus granulosus sensu lato*, which causes cystic echinococcosis (CE), or *E. multilocularis*, which causes alveolar echinococcosis (AE). In China, a national survey showed that CE is endemic in at least 368 counties in northwest China and is co-endemic with AE in 115 of those counties. The role of dogs in transmitting echinococcosis in these areas is significant, and efforts to dose dogs with monthly [deworming](#) treatments have been difficult to implement.

In the new work, Xiao-Nong Zhou of the Chinese Center for Disease Control and Prevention and colleagues developed a deworming collar which delivers praziquantel (PZQ), known to be the most effective deworming drug. They designed the collar to be waterproof, anti-collision, cold-proof, and to automatically deliver a regular, quantitative dose of PZQ. 18 smart deworming collars were used for [field testing](#) in Seni district of China, and 523 collars were tested in Henzuo city. Dogs for the trial were randomly selected from all registered dogs in each jurisdiction.

In pre-[field trials](#), the 551 collars were 100.0% anti-collision, 99.5% waterproof and 100.0% coldproof, and the automatic PZQ delivery occurred 87.8% of the time, even with the collars being attached to dogs for 12 months in the harsh climates of remote locations on the Tibetan Plateau. The compliance rate of dog owners to attach the collar to their dog was 94.7% in Seni district and 88.8% in Hezuo city. When compared to a control group, dogs in Seni district wearing the smart collar had a 0.182 times risk of a positive Echinococcus antigen test (95%CI 0.049-0.684, P=0.012) and dogs in Hezuo had a 0.336 times risk of a positive antigen test (95% CI 0.178-0.706, P=0.003).

"In order to prevent the transmission of echinococcosis from dogs to humans and livestock completely, we developed a smart Internet of Things (IoT)-based deworming collar which can deliver PZQ baits for [dogs](#) automatically and regularly," the researchers say. "Two pilot studies have showed that it is an excellent alternative to existing manual deworming methods, and the difficulties associated with performing deworming in remote areas with scarce resources can be overcome."

More information: Yang S-J, Xiao N, Li J-Z, Feng Y, Ma J-Y, Quzhen G-S, et al. (2021) Smart deworming collar: A novel tool for reducing Echinococcus infection in dogs. *PLoS Negl Trop Dis* 15(7): e0009443. doi.org/10.1371/journal.pntd.0009443

Provided by Public Library of Science

Citation: 'Smart collar' could prevent tapeworms in dogs (2021, July 8) retrieved 26 April 2024 from <https://phys.org/news/2021-07-smart-collar-tapeworms-dogs.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is

provided for information purposes only.