


Scientists analyze how the pandemic affected air quality

July 21 2021, by Sofie Bates



Drops in NO₂ during the pandemic were unequal for the most white (blue dots) and least white (orange dots) neighborhoods in the largest metropolitan areas. The solid lines indicate pre-pandemic NO₂ levels and the dashed lines NO₂ levels during the height of pandemic-related lockdowns. Credit: Kerr et al. 2021

The coronavirus pandemic has changed the way we live, and several studies have documented how those widespread changes in human behavior have impacted the environment. NASA scientists and others using data from NASA and our partner satellites have shown that air pollution levels dropped significantly during COVID-19. A new, NASA-funded study, conducted by scientists at The George Washington University (GW) in Washington, D.C., zoomed in on the 15 largest metropolitan areas in the United States to see how the drop in air pollution differed from neighborhood to neighborhood. The paper was published July 20 in the journal *Proceedings of the National Academy of Sciences*.

We talked to the lead author on the study, Gaige Kerr, about how the COVID-19 pandemic led to better air quality—and how those improvements were unequal for people of different races, ethnicities and socioeconomic levels. Kerr is a research scientist at GW. The interview has been lightly edited for clarity.

What is nitrogen dioxide and how is it related to air quality?

Nitrogen dioxide, or NO_2 , is a trace gas in the atmosphere. It's one of the six air pollutants regulated by the U.S. Environmental Protection Agency (EPA) under the Clean Air Act. Even though NO_2 is only present in small amounts, it's very harmful for human health and can trigger respiratory illnesses like asthma. NO_2 also leads to the formation of ozone near Earth's surface, another harmful air pollutant.

NO_2 can come from natural things like lightning or microbes in the soil. But in cities, the majority of NO_2 stems from human activity and fossil fuel combustion. Roughly 50% of the ambient NO_2 comes from traffic emissions. The other large sources come from power plants, incinerators

and factories.

What happened with NO₂ levels when the COVID-19 pandemic triggered lockdowns in March 2020?


During COVID, we had the opportunity to see how taking many cars off of the road and planes out of the skies affected air pollution in the real world in this unintended experiment. We know from past research that there are NO₂ pollution disparities based on several factors—most notably race, ethnicity and income—and that communities of color and lower socioeconomic status face much higher concentrations of NO₂. So, we wanted to understand how this unprecedented, extraordinary drop in human activity and emissions impacted NO₂ disparities.

In cities, NO₂ levels plummeted at first. That had a lot to do with the drop in traffic and travel, since vehicle traffic is the largest contributor to NO₂ in cities. We saw widespread decreases in NO₂ across urban areas in the United States during the pandemic, but the magnitude varied. NO₂ levels dropped by about 10% to 35% on average, depending on the [city](#). New York City and Los Angeles had very large drops, but NO₂ disparities across different racial, ethnic and socioeconomic groups were very large in these cities.

How did you determine NO₂ levels for different neighborhoods?

We used data from the TROPOspheric Monitoring Instrument (TROPOMI) on the European Commission's Copernicus Sentinel-5P satellite. With TROPOMI, in near real time, we can look at NO₂ levels in the atmosphere at a very high resolution—like neighborhood by neighborhood. We compared TROPOMI's NO₂ measurements with data from the U.S. Census Bureau from the American Community Survey. We collected all of the satellite measurements within a census tract—which we can think of as the size of a small neighborhood—and

paired that with the demographic data like race and ethnicity, median household income, education level and vehicle ownership.



Comparing maps of NO₂ levels, median household income and whiteness shows that air quality improvements were unequal between groups of people. In the map of NO₂ level compared to the average for NYC (left), blue indicates a larger drop in NO₂. That area overlaps with lower median household income (center) and a lower percentage of white residents (right). Credit: Kerr et al. 2021


What trends did you see when you looked at how the drop in NO₂ differed between neighborhoods?

The largest decreases in NO₂ level occurred in the metropolitan area neighborhoods with a larger non-white population. But even with these large reductions, the levels of NO₂ in those areas during the pandemic were still higher than the pre-pandemic levels of NO₂ in the neighborhoods with a higher percentage of white residents. In many of the cities we examined, there was no change in the size of NO₂ disparities between the most and least white or the highest or lowest income neighborhoods, despite an overall decrease in NO₂.

Your team analyzed NO₂ levels in the 15 largest metropolitan areas in the United States. Did you see this result in all of them?

Yes, we saw widespread decreases in NO₂ across urban areas in the United States during the pandemic, but the magnitude varied. NO₂ levels dropped by roughly 10% to 35% depending on the city.

We also did a deep dive into Detroit, New York, and Atlanta by looking at maps of NO₂ data and overlaying that with information from the U.S. Census Bureau. For example, in Detroit, we found the largest NO₂ drops were along the Detroit River. When we looked at what pollution sources lie around these neighborhoods, we saw that they were boxed in by a very busy interstate on one side and, on the other side, one of the busiest border crossings in North America, from Detroit to Windsor, Ontario, in Canada. There's usually a lot of heavy-duty trucks idling at the border crossing, waiting for their turn to get through customs and border patrol. So, it makes sense why we saw some of the largest NO₂ drops in those neighborhoods in Detroit.



The magnitude of the NO₂ drop that different groups experienced during the pandemic varied greatly by income and education level, racial and ethnic background, and household vehicle ownership. Credit: Kerr et al. 2021

In New York City, we found that the largest decrease in NO₂ pollution occurred in Harlem and the Bronx. This part of the city has been referred to as "asthma alley" due to asthma rates well above the national and state average. These neighborhoods and their citizens, who are primarily Black and Hispanic, face a barrage of pollution from heavy-duty trucking and industry.

In Atlanta, the largest NO₂ drops were in the southwest part of the city where there's a large international airport. That part of the city is also home to a majority black and low-income population. In addition to all of the highways and interstates in that part of the city, it's likely that emissions related to aviation are contributing to the NO₂ drops. The Guardian reported that in March of 2020 around 50% of flights had been canceled from Atlanta. So, there was not only a lot less vehicle traffic to the airport but also fewer planes flying in and out of the airport.

So, even with the improvements in air quality during the pandemic, NO₂ levels for communities of color were still higher than the pre-pandemic NO₂ levels for the whitest communities. What might be driving that discrepancy?

When we looked at the distribution of major roads in relation to each census tract, we found that the areas with the biggest drops in NO₂ had about nine to 10 times more highways and interstates nearby than the census tracts with the smallest drops. The neighborhoods with a larger non-white population have about five or six times more highways and interstates nearby than the most-white [neighborhoods](#).

Something else I thought was really interesting is that we found that census tracts with very low vehicle ownership in low-income communities and communities of color had some of the largest NO₂ drops during the lockdown. This suggests that some of the [air pollution](#)

experienced by people living in these areas is not caused by their own consumption of fossil fuels that produce air pollutants like NO₂.

More information: Gaige Hunter Kerr et al, COVID-19 pandemic reveals persistent disparities in nitrogen dioxide pollution, *Proceedings of the National Academy of Sciences* (2021). [DOI: 10.1073/pnas.2022409118](https://doi.org/10.1073/pnas.2022409118)

Provided by NASA

Citation: Scientists analyze how the pandemic affected air quality (2021, July 21) retrieved 20 September 2024 from <https://phys.org/news/2021-07-scientists-pandemic-affected-air-quality.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.