


Layered graphene with a twist displays unique quantum confinement in 2D

23 August 2021



Staff scientist Jurek Sadowski (left) and postdoc Zhongwei Dai at the Quantum Material Press (QPress) facility at the Center for Functional Nanomaterials (CFN) at Brookhaven National Laboratory. The large circular piece is the central QPress robot, with various modules attached on the sides for sample annealing, film deposition, plasma cleaning, and sample libraries. The full QPress system, still under development, will automate the stacking of 2-D materials into layered structures with exotic properties for quantum applications. Credit: Brookhaven National Laboratory

Scientists studying two different configurations of bilayer graphene—the two-dimensional (2D), atom-thin form of carbon—have detected electronic and optical interlayer resonances. In these resonant states, electrons bounce back and forth between the two atomic planes in the 2D interface at the same frequency. By characterizing these states, they found that twisting one of the graphene layers by 30 degrees relative to the other, instead of stacking the layers directly on top of each other, shifts the resonance to a lower energy. From this result, just published in *Physical Review Letters*, they deduced that the distance between the two layers increased significantly in the twisted configuration, compared to the stacked one. When this distance changes, so do the interlayer

interactions, influencing how electrons move in the bilayer system. An understanding of this electron motion could inform the design of future quantum technologies for more powerful computing and more secure communication.

"Today's computer chips are based on our knowledge of how electrons move in semiconductors, specifically silicon," said first and co-corresponding author Zhongwei Dai, a postdoc in the Interface Science and Catalysis Group at the Center for Functional Nanomaterials (CFN) at the U.S. Department of Energy (DOE)'s Brookhaven National Laboratory. "But the physical properties of silicon are reaching a physical limit in terms of how small transistors can be made and how many can fit on a chip. If we can understand how electrons move at the small scale of a few nanometers in the reduced dimensions of 2D materials, we may be able to unlock another way to utilize electrons for quantum information science."


At a few nanometers, or billionths of a meter, the size of a material system is comparable to that of the wavelength of electrons. When electrons are confined in a space with dimensions of their wavelength, the material's electronic and optical properties change. These quantum confinement effects are the result of quantum mechanical wave-like motion rather than classical mechanical motion, in which electrons move through a material and are scattered by random defects.

For this research, the team selected a simple material model—graphene—to investigate quantum confinement effects, applying two different probes: electrons and photons (particles of light). To probe both electronic and optical resonances, they used a special substrate onto which the graphene could be transferred. Co-corresponding author and CFN Interface Science and Catalysis Group scientist Jurek Sadowski had previously designed this substrate for the Quantum Material Press (QPress). The QPress is an automated tool under

development in the CFN Materials Synthesis and Characterization Facility for the synthesis, processing, and characterization of layered 2D materials. Conventionally, scientists exfoliate 2D material "flakes" from 3D parent crystals (e.g., graphene from graphite) on a silicon dioxide substrate several hundred nanometers thick. However, this substrate is insulating, and thus electron-based interrogation techniques don't work. So, Sadowski and CFN scientist Chang-Yong Nam and Stony Brook University graduate student Ashwanth Subramanian deposited a conductive layer of titanium oxide only three nanometers thick on the silicon dioxide substrate.

"This layer is transparent enough for optical characterization and determination of the thickness of exfoliated flakes and stacked monolayers while conductive enough for electron microscopy or synchrotron-based spectroscopy techniques," explained Sadowski.

In the Charlie Johnson Group at the University of Pennsylvania—Rebecca W. Bushnell Professor of Physics and Astronomy Charlie Johnson, postdoc Qicheng Zhang, and former postdoc Zhaoli Gao (now an assistant professor at the Chinese University of Hong Kong)—grew the graphene on metal foils and transferred it onto the titanium oxide/silicon dioxide substrate. When graphene is grown in this way, all three domains (single layer, stacked, and twisted) are present.



(a) Schematics of the experimental setup for electron and photon scattering. (b) An atomic model of the pattern formed by the twisted bilayer graphene (30°-tBLG) crystal structure. (c) A low-energy electron microscope image of a typical sample area containing 30°-tBLG, stacked bilayer graphene (AB-BLG), and single-layer graphene (SLG). (d) A low-energy electron diffraction pattern on a 30°-tBLG area. Credit: Brookhaven National Laboratory

Then, Dai and Sadowski designed and carried out experiments in which they shot electrons into the material with a low-energy electron microscope (LEEM) and detected the reflected electrons. They also fired photons from a laser-based optical microscope with a spectrometer into the material and analyzed the spectrum of light scattered back. This confocal Raman microscope is part of the QPress cataloger, which together with image-analysis software, can pinpoint the locations of sample areas of interest.

"The QPress Raman microscope enabled us to quickly identify the target sample area, accelerating our research," said Dai.

Their results suggested that the spacing between layers in the twisted graphene configuration increased by about six percent relative to the non-twisted configuration. Calculations by theorists at the University of New Hampshire verified the unique resonant electronic behavior in the twisted configuration.

"Devices made out of rotated graphene may have very interesting and unexpected properties because of the increased interlayer spacing in which electrons can move," said Sadowski.

Next, the team will fabricate devices with the twisted graphene. The team will also build upon initial experiments conducted by CFN staff scientist Samuel Tenney and CFN postdocs Calley Eads and Nikhil Tiwale to explore how adding different materials to the layered structure impacts its electronic and optical properties.

"In this initial research, we picked the simplest 2D material system we can synthesize and control to understand how electrons behave," said Dai. "We plan to continue these types of fundamental studies, hopefully shedding light on how to manipulate materials for quantum computing and communications."

More information: Zhongwei Dai et al, Quantum-Well Bound States in Graphene Heterostructure Interfaces, *Physical Review Letters* (2021). DOI: [10.1103/PhysRevLett.127.086805](https://doi.org/10.1103/PhysRevLett.127.086805)

Provided by Brookhaven National Laboratory

APA citation: Layered graphene with a twist displays unique quantum confinement in 2D (2021, August 23) retrieved 27 October 2021 from <https://phys.org/news/2021-08-layered-graphene-unique-quantum-confinement.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.