


Moth wingtips an 'acoustic decoy' to thwart bat attack, scientists find

9 September 2021



The Atlas moth (*Attacus atlas*) has a strong anti-bat acoustic decoy at the tip of its forewings. Composite image with photograph on right half and acoustic tomography on the left. Colour indicates echo strength on a dB scale and red indicates highest echo amplitude. Note the red highly reflecting stripe created by the rippled part of the wingtip. Credit: T. Neil and M Holderied

Wingtips of certain species of silkworm are structured to reflect sound and throw off attackers, according to a new study.

Researchers at the University of Bristol have discovered that the tips of some saturniid moth forewings are curiously rippled and folded. They found that these unique structures strongly reflect sound, meaning that a bat hunting using echolocation is more likely to attack the wingtip region of the moth over the body, potentially saving the moth's life.

They also discovered that the ripples and folds of the forewing tips have evolved to act as hemispheric and corner retroreflectors respectively, meaning that they reflect sound strongly back to its point of origin. Coupled together, the folds and ripples of these wingtips cover a huge range of incident sounds angles,

meaning that over the entire wingbeat cycle of a flying moth and most possible positions of an attacking bat, the wingtip would consistently produce the strongest echoes. The acoustic protection of wingtips is even stronger than that of common hindwing decoys.

Prof Marc Holderied of Bristol's School of Biological Sciences explained: "We have demonstrated that the folded and rippled wingtips on the forewings of some silkworms act as acoustic decoys.

"Structurally, the wingtips act as acoustic retroreflectors, reflecting sound back to its source from numerous angles, meaning a bat would be more likely to strike the wingtip over the more vulnerable body of the moth."



Photograph of the Atlas moth (*Attacus atlas*). This large silkworm has the strongest known wingtip decoy with ripples and folds. Credit: T. Neil

The findings, published today in *Current Biology*, are the latest revelation in the bat-moth acoustic arms race—the battle between bats which

hunt moths using echolocation, and the subsequent evolution of different defensive strategies amongst moths to increase their chances of survival.

Towed acoustic decoys are a well-established defense amongst some silkmoths. These species have evolved elongated hindwings which terminate in a coiled and twisted end. The morphology of these elongated hindwings means that they generate very strong echoes, so much so that they will often divert a bat's acoustic gaze towards them, away from the exposed body of the moth, causing the bat to strike the expendable tail of the moth or miss the [moth](#) all together.

Lead author Dr. Thomas Neil said: "There are many silkmoths that do not have these elongated hindwings, and we were interested in how they might protect themselves from bats. Through our research we discovered that there are many silkmoths that have rippled and folded structures not on the tips of their elongated hindwings but on the tips of their forewings. These resembled the twisted hindwing structures seen in other moths and so we wanted to know if they might also serve as an acoustic decoy to thwart a bat's attack.

"To test this theory, we used innovative acoustic tomography analysis. We recorded echoes from moths from over 10,000 angles, to compare whether the echoes coming from the wingtips of these moths were stronger than the echoes from the body. If the echoes coming from the rippled and folded wingtips were stronger than that of the body, this would indicate that they were indeed acoustic decoys.

"Conclusive support for the idea that the forewing reflector is an acoustic decoy comes from our finding that acoustic forewing decoys always evolved as an alternative to acoustic hindwing decoys, with there being no species known to possess both."



Photograph of the Chinese tussar moth (*Antheraea pernyi*). This silkmoth has a strong wingtip decoy based on ripples. Credit: T. Neil

Now the researchers will try and collect behavioral data to corroborate their findings in the lab. They plan to monitor bats and moths with varying levels of folded wing morphologies to see how much of a survival advantage it really gives them.

Prof Holderied added: "The results of this study introduce another exciting aspect to the story of the bat-moths acoustic arms race. We have identified a novel form of acoustic defense amongst silkmoths which may give them an advantage over hunting bats. Wider implications might include improved man-made anti radar and sonar decoy architectures."

"Wingtip folds and ripples on saturniid moths create decoy echoes against bat biosonar" is published in *Current Biology* by Thomas Neil, Ella Kennedy, Brogan Harris, and Marc Holderied.

More information: Wingtip folds and ripples on saturniid moths create decoy echoes against bat biosonar, *Current Biology* (2021).

Provided by University of Bristol

APA citation: Moth wingtips an 'acoustic decoy' to thwart bat attack, scientists find (2021, September 9) retrieved 2 December 2021 from <https://phys.org/news/2021-09-moth-wingtips-acoustic-decoy-thwart.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.