


How do higher waves cause more ice clouds? Research expedition into arctic sea explains

September 17 2021



Oceanic supply of ice-nucleating particles and its effect on ice cloud formation: A case study in the Arctic Ocean during a cold-air outbreak in early winter
Inoue et al. (2021) | *Geophysical Research Letters* | DOI: 10.1029/2021GL094646

Global warming is causing a rapid decline in sea-ice area, which affects weather patterns and, surprisingly, increases wave height in the Arctic. In a new study published in *Geophysical Research Letters*(10.1029/2021GL094646), Japanese scientists analyzed data from a 2018 research expedition into Chukchi Sea to demonstrate the peculiar link that exists between sea spray induced by high waves and the formation of ice-containing clouds. Their results pave the way for more accurate climate change and sea-ice models. Credit: NIPR/JAMSTEC/Kitami Institute of Technology

Some of the most well-known and feared aspects of climate change are its potential effects on weather patterns and how this could accelerate the melting of natural ice. Research has already proved that the area of sea-ice in the Arctic is rapidly declining due to global warming, and that temperature and moisture content across the Arctic have changed substantially. Unfortunately, understanding exactly how these changes affect cloud formation in the region is very challenging, and cloud composition and phase are important aspects to consider in predictive numerical models.

In a recent study published in *Geophysical Research Letters*, a team of scientists led by Dr. Jun Inoue of the National Institute of Polar Research, Japan, sought to answer a peculiar question: can higher waves in the Arctic Sea promote the development of ice-containing clouds? This question may seem strange at first, because most people would not have fathomed that a link could exist between those two natural phenomena. However, as the findings of this study indicate, it's likely that there is one.

The field data used in the study was gathered in November 2018 during an expedition to the Chukchi Sea in the Arctic region aboard RV Mirai, a Japanese research vessel. Previous studies in the area had revealed that the sea-ice decline in the Arctic led to more frequent active weather systems, stronger winds, and taller waves. The research team suspected that these factors could affect cloud formation and composition because crashing waves and strong winds can cause organic particles on the sea surface to become dispersed through the atmosphere in the form of sea spray. Once these suspended [organic particles](#) reach an altitude high enough, they act as "seeds" that facilitate the formation of ice crystals, earning them the name of "ice-nucleating particles" (INPs). These ice crystals keep growing by freezing the surrounding water droplets, thereby forming what's known as ice clouds.

To prove this hypothesis, Dr. Inoue and his crew on the RV Mirai periodically deployed various measurement instruments at key locations in the Chukchi Sea over the course of 12 days. Cloud particle sensors were balloon-launched from the ship to analyze the phase of clouds, ambient aerosols were regularly sampled on board for chemical analysis, and wave height and wind speed measurements were constantly made. Moreover, the researchers conducted turbidity measurements from different depths to clarify the relationship between weather and oceanic conditions.

After analyzing all the gathered data, the scientists managed to paint a clearer, evidence-supported view of the situation. "Chukchi Sea is relatively shallow, with a mean depth of only 40 meters. There, a mixed ocean layers develops and taps into the seafloor, which cloud provide a reservoir of INPs that get lifted by turbulent kinetic energy," explains Dr. Inoue, "Sea spray induced by strong winds and high waves brings these INPs to the atmosphere, promoting the formation of ice clouds." He adds that this is one of the first papers to simultaneously link oceanic structure, sea surface conditions, and aerosol and cloud characteristics.

The insight gained from this study is very important if we are to accurately predict the effects of global warming on the Arctic. Ice clouds reflect much less shortwave solar radiation than water clouds, and thus the phase of [clouds](#) greatly affects the surface heat budget of the polar regions. They may also increase the amount of snowfall, which in turn positively affects sea-ice formation. "Understanding the relationship between [cloud formation](#) and the new sea state originating from the recent Arctic sea-ice decline is critical for skillful weather and sea-ice forecasts, as well as future climate projections," highlights Dr. Inoue. Let us hope further studies in the Arctic allow us to elucidate all the fine details and hidden interactions that dictate the weather so that the consequences of climate change don't catch us off-guard.

More information: Jun Inoue et al, Oceanic Supply of Ice-Nucleating Particles and Its Effect on Ice Cloud Formation: A Case Study in the Arctic Ocean During a Cold-Air Outbreak in Early Winter, *Geophysical Research Letters* (2021). [DOI: 10.1029/2021GL094646](https://doi.org/10.1029/2021GL094646)

Provided by Research Organization of Information and Systems

Citation: How do higher waves cause more ice clouds? Research expedition into arctic sea explains (2021, September 17) retrieved 19 September 2024 from <https://phys.org/news/2021-09-higher-ice-clouds-arctic-sea.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.